

Recombinant Human TRPV1 293 Cell Lysate

Cat. No. TRPV1-735HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for transient receptor potential cation channel, subfamily V, member 1 (TRPV1), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	TRPV1 transient receptor potential cation channel, subfamily V, member 1 [Homo sapiens]
Official Symbol	TRPV1
Synonyms	TRPV1; transient receptor potential cation channel, subfamily V, member 1; vanilloid receptor subtype 1 , VR1; transient receptor potential cation channel subfamily V member 1; OTRPC1; capsaicin receptor; osm-9-like TRP channel 1; vanilloid receptor subtype 1; transient receptor potential vanilloid 1a; transient receptor potential vanilloid 1b; VR1; DKFZp434K0220;
Gene ID	7442
mRNA Refseq	NM_018727
Protein Refseq	NP_061197
MIM	602076
UniProt ID	Q8NER1
Chromosome Location	17p13.3
Pathway	Neuroactive ligand-receptor interaction, organism-specific biosystem; Neuroactive ligand-receptor interaction, conserved biosystem; Trk receptor signaling mediated by PI3K and PLC-gamma, organism-specific biosystem; Trk receptor signaling mediated

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



by the MAPK pathway, organism-specific biosystem;

☎ Tel: 1-631-559-9269 1-516-512-3133

☎ Email: info@creative-biomart.com ☎ Fax: 1-631-938-8127

☎ 45-1 Ramsey Road, Shirley, NY 11967, USA