

Active Recombinant Mouse Tat protein, His-tagged

Cat. No. Tat-2115M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Mouse Tat protein(Q8QZR1)(190-454 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Mouse
Source	E.coli
ProteinLength	190-454 aa
Form	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Bio-activity	<p>Tyrosine aminotransferase (TAT) is an enzyme present in the liver and catalyzes the conversion of tyrosine to 4-hydroxyphenylpyruvate. In humans, the tyrosine aminotransferase protein is encoded by the TAT gene. A deficiency of the enzyme in humans can result in what is known as Type II Tyrosinemia, wherein there is an abundance of tyrosine as a result of tyrosine failing to undergo an aminotransferase reaction to form 4-hydroxyphenylpyruvate. Besides, Glutamine synthetase (GS) has been identified as an interactor of TAT, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse TAT and recombinant mouse GS. Briefly, TAT were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to GS-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-TATpAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µ stop solution to the wells and read at 450nm immediately. The binding activity of TAT and GS was shown, and this effect was in a dose dependent manner.

Molecular Mass	33.7kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 95%
Storage	Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	Tat tyrosine aminotransferase [Mus musculus]
Official Symbol	Tat
Synonyms	TAT; tyrosine aminotransferase; tyrosine transaminase; L-tyrosine:2-oxoglutarate aminotransferase; MGC37772; MGC37789; MGC37790; MGC37819; MGC37828; MGC37842;
Gene ID	234724
mRNA Refseq	NM_146214

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Protein Refseq

NP_666326

SDS-PAGE



Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA