

Recombinant Mouse Tnfsf13b protein, His/S-tagged

Cat. No. Tnfsf13b-220R Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant Mouse Tnfsf13b fused with His/S tag was expressed in E. coli.

Species Mouse

Source E.coli

Description

B-cell activating factor, also known as BAFF, TALL-1, TNAK, and zTNF4, is a member of the TNF ligand superfamily designated TNFSF13B. Produced by macrophages, dendritic cells, and T lymphocytes, BAFF promotes the survival of B cells and is essential for B cell maturation. BAFF binds to three TNF receptor superfamily members: B-cell maturation antigen (BCMA/TNFRSF17), transmembrane activator and calcium-modulator and cyclophilin ligand interactor (TACI/TNFRSF13B) and BAFF receptor (BAFF R/BR3/TNFRSF 13C). These receptors are type III transmembrane proteins lacking a signal peptide. Whereas TACI and BCMA bind BAFF and another TNF superfamily ligand, APRIL (a proliferation-inducing ligand), BAFF R selectively binds BAFF. The BAFF R extracellular domain lacks the TNF receptor canonical cysteine-rich domain (CRD) and contains only a partial CRD with four cysteine residues. Human and mouse BAFF R share 56% aa sequence identity. BAFF R is highly expressed in spleen, lymph node and resting B cells. It is also expressed at lower levels in activated B cell, resting CD4+ T cells, thymus and peripheral blood leukocytes.

Form Lyophilized from sterile PBS, pH 7.4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity > 95 % as determined by SDS-PAGE

Storage Store at -70 centigrade. Avoid repeated freeze/thaw cycles.

GENE INFORMATION

Gene Name [Tnfsf13b tumor necrosis factor \(ligand\) superfamily, member 13b \[Mus musculus \]](#)

Official Symbol [Tnfsf13b](#)

Synonyms TNFSF13B; tumor necrosis factor (ligand) superfamily, member 13b; tumor necrosis factor ligand superfamily member 13B; B-cell-activating factor; b cell-activating factor; BAFF; BLyS; TALL1; THANK; zTNF4; TALL-1; TNFSF20; D8Erttd387e; MGC124060; MGC124061;

Gene ID [24099](#)

mRNA Refseq [NM_033622](#)

Protein Refseq [NP_296371](#)

UniProt ID [Q9WU72](#)

Chromosome Location 8 A1.1; 8 4.55 cM

Pathway Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Intestinal immune network for IgA production, organism-specific biosystem; Intestinal immune network for IgA production, conserved biosystem; Rheumatoid arthritis, organism-specific biosystem; Rheumatoid arthritis, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

cytokine activity; tumor necrosis factor receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA