

## Recombinant Human UBA1 cell lysate

Cat. No. UBA1-420HCL Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Human UBE1 / UBA1 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
<b>Species</b>	Human
<b>Source</b>	Insect Cells
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

## GENE INFORMATION

**Gene Name**

[UBA1 ubiquitin-like modifier activating enzyme 1 \[ Homo sapiens \]](#)

**Official Symbol**

UBA1

**Synonyms**

UBA1; ubiquitin-like modifier activating enzyme 1; A1S9T, GXP1, UBE1, ubiquitin activating enzyme E1 , ubiquitin activating enzyme E1 (A1S9T and BN75 temperature sensitivity complementing); ubiquitin-like modifier-activating enzyme 1; POC20; POC20 centriolar protein homolog (Chlamydomonas); ubiquitin activating enzyme E1 homolog (yeast); UBE1X; POC20 centriolar protein homolog; UBA1, ubiquitin-activating enzyme E1 homolog A; A1S9T and BN75 temperature sensitivity complementing; A1S9; A1ST; GXP1; UBE1; A1S9T; AMCX1; SMAX2; UBA1A; MGC4781;

**Gene ID**

[7317](#)

**mRNA Refseq**

[NM\\_003334](#)

**Protein Refseq**

[NP\\_003325](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>MIM</b>	314370
<b>UniProt ID</b>	P22314
<b>Chromosome Location</b>	Xp11.23
<b>Pathway</b>	Adaptive Immune System, organism-specific biosystem; Antigen processing: Ubiquitination and Proteasome degradation, organism-specific biosystem; Class I MHC mediated antigen processing & presentation, organism-specific biosystem; Immune System, organism-specific biosystem; Parkinsons disease, organism-specific biosystem;
<b>Function</b>	ATP binding; ligase activity; nucleotide binding; protein binding; small protein activating enzyme activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA