

## Active Recombinant Human USP28, GST-tagged

Cat. No. USP28-188H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant human USP28 (amino acid residues 1-1077), fused with N-terminal GST, was expressed in E.coli.
<b>Species</b>	Human
<b>Source</b>	E.coli
<b>ProteinLength</b>	1-1077 a.a.
<b>Description</b>	Ubiquitin carboxyl-terminal hydrolase 28 (Ubiquitin Specific Protease 28; USP28) is a member of the cysteine protease enzyme family and cloning of the human gene was first described by Nagase et al. (2000). USP28 shares 51.4% identity with USP25 and is highly expressed in the heart and skeletal muscle (Valero et al., 2001). Both USP28 and Checkpoint kinase Chk2 are required for DNA-damage-induced apoptosis, and they accomplish this partly through regulation of the p53 induction of proapoptotic genes. USP28 is required to stabilize Chk2, p53 binding protein 53BP1 and claspin in response to DNA double strand breaks (Zhang et al., 2006).
<b>Form</b>	50 mM HEPES pH 7.5, 150 mM sodium chloride, 2 mM dithiothreitol, 10% glycerol
<b>Bio-activity</b>	Deubiquitylase Enzyme Assay: The activity of GST-USP28 was validated by determining the increase in fluorescence measured as a result of the enzyme catalysed cleavage of the fluorogenic substrate Ubiquitin-Rhodamine110-Glycine generating Ubiquitin and Rhodamine110-Glycine. Incubation of the substrate in the presence or absence of GST-USP28 was compared confirming the deubiquitylating

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	activity of GST-USP28.
<b>Molecular Mass</b>	~149 kDa
<b>Purity</b>	>70% by SDS-PAGE
<b>Storage</b>	12 months at -70°C. Avoid multiple freeze/thaw cycles.
<b>Concentration</b>	0.5 mg/ml

## GENE INFORMATION

<b>Gene Name</b>	USP28 ubiquitin specific peptidase 28 [ Homo sapiens ]
<b>Official Symbol</b>	USP28
<b>Synonyms</b>	USP28; ubiquitin specific peptidase 28; ubiquitin specific protease 28; ubiquitin carboxyl-terminal hydrolase 28; KIAA1515; ubiquitin thioesterase 28; deubiquitinating enzyme 28; ubiquitin thiolesterase 28; ubiquitin-specific-processing protease 28;
<b>Gene ID</b>	<a href="#">57646</a>
<b>mRNA Refseq</b>	<a href="#">NM_020886</a>
<b>Protein Refseq</b>	<a href="#">NP_065937</a>
<b>MIM</b>	<a href="#">610748</a>
<b>UniProt ID</b>	<a href="#">Q96RU2</a>
<b>Chromosome Location</b>	11q23

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



**Function**

cysteine-type peptidase activity; peptidase activity; protein binding; ubiquitin thiolesterase activity; ubiquitin thiolesterase activity; ubiquitin-specific protease activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA