

Recombinant Human USP28 cell lysate

Cat. No. USP28-1895HCL Lot. No. (See product label)

SPECIFICATION

Species

Human

Description

The ubiquitin-dependent protein degradation pathway is essential for proteolysis of intracellular proteins and peptides. Enzymes that remove ubiquitin from ubiquitin-conjugated peptides, like USP28, affect the fate and degradation of intracellular proteins and are essential for maintenance of cell-free ubiquitin pools (Valero et al., 2001).

Size

100 ul

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Applications

Western Blot;

GENE INFORMATION

Gene Name

USP28 ubiquitin specific peptidase 28 [Homo sapiens]

Official Symbol

USP28

Synonyms

USP28; ubiquitin specific peptidase 28; ubiquitin specific protease 28; ubiquitin carboxyl-terminal hydrolase 28; KIAA1515; ubiquitin thioesterase 28; deubiquitinating enzyme 28; ubiquitin thiolesterase 28; ubiquitin-specific-processing protease 28;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Gene ID	57646
mRNA Refseq	NM_020886
Protein Refseq	NP_065937
MIM	610748
UniProt ID	Q96RU2
Chromosome Location	11q23
Function	cysteine-type peptidase activity; peptidase activity; protein binding; ubiquitin thiolesterase activity; ubiquitin thiolesterase activity; ubiquitin-specific protease activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA