

Active Recombinant Human VCAM1 protein, His-tagged

Cat. No. VCAM1-3371H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human VCAM1 protein(P19320)(109~318 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	109-318 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Vascular cell adhesion protein 1 (VCAM-1) is a cell adhesion molecule and a member of the immunoglobulin superfamily. Alternatively spliced forms are known to occur, but the most common form is a type I transmembrane protein with a 674 amino acid extracellular segment that includes seven C2-type immunoglobulin domains. VCAM-1 is expressed by activated endothelial cells and certain other cell types including macrophages, dendritic cells, neurons, smooth muscle cells, fibroblasts, and oocytes. Besides, Integrin Alpha 4 (ITGa4) has been identified as an interactor of VCAM-1, thus a binding ELISA assay was conducted to detect the interaction of recombinant human VCAM-1 and recombinant human ITGa4. Briefly, biotin-linked VCAM-1 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to ITGa4-coated microtiter wells and incubated for 1h at 37°C. Wells were washed with PBST 3 times and incubation with Streptavidin-HRP for 30min,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

then wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 µstop solution to the wells and read at 450nm immediately. The binding activity of recombinant human VCAM-1 and recombinant human ITGa4 was shown, the EC50 for this effect is 0.261 ug/mL.

Molecular Mass	25.0kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 90% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	VCAM1 vascular cell adhesion molecule 1 [Homo sapiens]
Official Symbol	VCAM1
Synonyms	VCAM1; vascular cell adhesion molecule 1; vascular cell adhesion protein 1; CD106; CD106 antigen; INCAM-100; MGC99561; DKFZp779G2333;
Gene ID	7412
mRNA Refseq	NM_001078

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_001069


MIM 192225

UniProt ID P19320

SDS-PAGE



Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA