

Recombinant Human VIM protein

Cat. No. VIM-233H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human VIM protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	465
Description	This gene encodes a type III intermediate filament protein. Intermediate filaments, along with microtubules and actin microfilaments, make up the cytoskeleton. The encoded protein is responsible for maintaining cell shape and integrity of the cytoplasm, and stabilizing cytoskeletal interactions. This protein is involved in neuritogenesis and cholesterol transport and functions as an organizer of a number of other critical proteins involved in cell attachment, migration, and signaling. Bacterial and viral pathogens have been shown to attach to this protein on the host cell surface. Mutations in this gene are associated with congenital cataracts in human patients.
Form	Lyophilized from a 0.2µm filtered solution in 30 % Acetonitrile and 0.1 % TFA.
Molecular Mass	Approximately 53.5 kDa, a single non-glycosylated polypeptide chain containing 465 amino acids.
AA Sequence	STRSVSSSSYRRMFGGPGTASRPSSRSYVTTSTRTYSLGSALRPSTSRSLYASSP GGVYATRSSAVRLRSSVPGVRLQDSVDFSLADAINTEFKNTRTNEKVELQELNDRF

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

ANYIDKVRFLEQQNKILLAELEQLKGQGKSRLGDLYEEEMRELRRQVDQLTNDKARV
 EVERDNLAEDIMRLREKLQEEMLQREEAENTLQSFQRQVDNASLARLDLERKVESL
 QEEIAFLKKLHEEEIQELQAQIQEQHVQIDVDVSKPDLTAALRDVRQQYESVAAKNLQ
 EAEWYKSKFADLSEAANRNNDALRQAKQESTEYRRQVQSLTCEVDALKGTNESL
 ERQMREMEENFAVEAANYQDTIGRLQDEIQNMKEEMARHLREYQDLLNVKMALDIE
 IATYRKLEGEESRISLPLPNFSSLNLRETNLDSLPLVDTHSKRTLLIKTVETRDGQVIN
 ETSQHDDLE

Endotoxin Less than 0.1 EU/g of rHuVimentin as determined by LAL method.

Purity >95% by SDS-PAGE and HPLC analysis.

Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.

Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 4 mM HCl to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name VIM

Official Symbol VIM

Synonyms VIM; vimentin; FLJ36605;

Gene ID 7431

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



mRNA Refseq	NM_003380
Protein Refseq	NP_003371
MIM	193060
UniProt ID	P08670

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA