

## Recombinant Human VTCN1 protein, His-tagged

Cat. No. VTCN1-662H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human VTCN1 (Phe29-Ala258) protein was fused to His-tag at C-terminus and expressed in human 293 cells (HEK293).
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	Phe29-Ala258
<b>Description</b>	V-set domain-containing T-cell activation inhibitor 1 (VTCN1) is also known as Immune costimulatory protein B7-H4, Protein B7S1, T-cell costimulatory molecule B7x, B7H4, which belongs to the immunoglobulin superfamily and BTN/MOG family. VTCN1 contains two Ig-like V-type (immunoglobulin-like) domains. The expression of VTCN1 is up-regulated by IL6 and IL10 and is inhibited by GM-CSF and IL4 on antigen-presenting cells (APCs). VTCN1/B7-H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. VTCN1 involved in promoting epithelial cell transformation.
<b>Predicted N Terminal</b>	Phe29
<b>Form</b>	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4, 10% trehalose.
<b>Molecular Mass</b>	The protein has a calculated MW of 26.1 kDa. The protein migrates as 43-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Endotoxin</b>	Less than 1.0 EU per µg by the LAL method.
<b>Purity</b>	>95% as determined by SDS-PAGE.
<b>Storage</b>	<p>For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower.</p> <p>Please avoid repeated freeze-thaw cycles.</p> <p>This product is stable after storage at:</p> <p>-20 to -70 centigrade for 12 months in lyophilized state;            -70 centigrade for 3 months under sterile conditions after reconstitution.</p>
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

## GENE INFORMATION

<b>Gene Name</b>	VTCN1
<b>Official Symbol</b>	VTCN1
<b>Synonyms</b>	VTCN1; V-set domain containing T cell activation inhibitor 1; V-set domain-containing T-cell activation inhibitor 1; B7 family member; H4; B7 superfamily member 1; B7 H4; B7H4; B7S1; B7X; FLJ22418; B7 family member, H4; T cell costimulatory molecule B7x; T-cell costimulatory molecule B7x; immune costimulatory protein B7-H4; B7-H4; B7h.5; VCTN1; PRO1291; RP11-229A19.4
<b>Gene ID</b>	79679
<b>mRNA Refseq</b>	NM_001253849
<b>Protein Refseq</b>	NP_001240778

 Tel: 1-631-559-9269 1-516-512-3133


 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**MIM** 608162


**UniProt ID** Q7Z7D3

**SDS-PAGE of  
VTCN1-662H**



Human B7-H4, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

**Bioactivity-ELISA of  
VTCN1-662H**




Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Immobilized Human B7-H4, His Tag at 2 µg/mL (100 µL/well) can bind Anti-B7-H4 MAb (Human IgG1) with a linear range of 0.2-6 ng/mL (QC tested).


**Bioactivity-ELISA**



Batch consistency of Bioactivity-ELISA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA