

Active Recombinant Human Wilms Tumor 1

Cat. No. WT1-1210H **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	Sf9 Cells
Description	<p>WT-1, the product of Wilms' Tumor suppressor gene <i>Wt1</i>, is a nuclear protein with structural motifs characteristic of transcription factors, including four C-terminal zinc fingers. While different pre-mRNA processing could result in 16 isoforms of the protein, the inclusion or exclusion of exon 5 and three amino acids (KTS) between zinc fingers 3 and 4 largely affects the activity of the WT-1 protein. Such a complex posttranscriptional regulation, particularly in splicing, may represent a major regulatory mechanism for tumorigenesis of the Wilms' tumor. With the inclusion of exon 5, WT-1 (+KTS) binds to both DNA and RNA and is RNase but not DNase sensitive. This form also colocalizes with splicing factor in a speckled nuclear particle, suggesting that WT-1 protein may function as both a transcription factor and a splicing regulator. The WT-1 protein (residue 1-449, including exon 5 and KTS) is expressed in baculovirus system and purified by an affinity column in combination with FPLC chromatography.</p>
Form	Liquid. Supplied in 20 mM Tris-HCl pH 8.0, 100 mM KCl, 0.2 mM EDTA, 1 mM DTT and 20% glycerol.
Purity	≥95% by SDS-PAGE.
Activity	1 ng is the amount sufficient for a gel mobility shift assay in a 20 µl reaction, 50 ng

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

are sufficient for reconstituted transcription assays and 100 ng are sufficient for a protein-protein interaction assay.

Usage For in vitro use only.

Storage Quality guaranteed for 12 months store at - 80°C. Avoid freeze / thaw cycles.

GENE INFORMATION

Gene Name [WT1 Wilms tumor 1 \[Homo sapiens \]](#)

Synonyms WT1; Wilms tumor 1; GUD; AWT1; WAGR; WT33; WIT-2; OTTHUMP00000037553; Wilms tumor protein

Gene ID [7490](#)

mRNA Refseq [NM_000378](#)

Protein Refseq [NP_000369](#)

MIM [607102](#)

UniProt ID [Q8IYZ5](#)

Chromosome Location 11p13


Function RNA binding; double-stranded DNA binding; metal ion binding; promoter binding; protein binding; transcription activator activity; transcription factor activity; zinc ion binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

PDB rendering based
on 1xf7.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA