

## Recombinant Human XRCC5 293 Cell Lysate

Cat. No. XRCC5-254HCL Lot. No. (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for X-ray repair complementing defective repair in Chinese hamster cells 5 (double-strand-break rejoining) (XRCC5) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	XRCC5 X-ray repair complementing defective repair in Chinese hamster cells 5 (double-strand-break rejoining) [ Homo sapiens ]
<b>Official Symbol</b>	XRCC5
<b>Synonyms</b>	XRCC5; X-ray repair complementing defective repair in Chinese hamster cells 5 (double-strand-break rejoining); X ray repair complementing defective repair in Chinese hamster cells 5 (double strand break rejoining; Ku autoantigen, 80kD); X-ray repair cross-complementing protein 5; KARP 1; Ku autoantigen; 80kDa; KU80; Ku86; KUB2; TLAA; CTC85; CTCBF; nuclear factor IV; Ku autoantigen, 80kDa; DNA repair protein XRCC5; thyroid-lupus autoantigen; 86 kDa subunit of Ku antigen; lupus Ku autoantigen protein p86; Ku86 autoantigen related protein 1; CTC box-binding factor 85 kDa subunit; ATP-dependent DNA helicase 2 subunit 2; ATP-dependent DNA helicase II 80 kDa subunit; NFIV; KARP1; KARP-1; FLJ39089;
<b>Gene ID</b>	7520
<b>mRNA Refseq</b>	NM_021141
<b>Protein Refseq</b>	NP_066964
<b>MIM</b>	194364
<b>UniProt ID</b>	P13010
<b>Chromosome Location</b>	2q35

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



**Pathway**

2-LTR circle formation, organism-specific biosystem; BARD1 signaling events, organism-specific biosystem; Coregulation of Androgen receptor activity, organism-specific biosystem; DNA Repair, organism-specific biosystem; DNA-PK complex, organism-specific biosystem; Disease, organism-specific biosystem; Double-Strand Break Repair, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA