

Active Recombinant Human TNFSF10 protein, His-tagged

Cat. No. TNFSF10-302H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human TNFSF10 protein(P50591)(114~281 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	114-281 aa
Form	Lyophilized from sterile 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Bio-activity	Protein TRAIL (TNF-Related Apoptosis-Inducing Ligand) is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. TRAIL induces apoptosis in transformed and tumor cells. This protein binds to several members of TNF receptor superfamily including TNFRSF10B/TRAILR2. Thus a binding ELISA assay was conducted to detect the interaction of TRAIL and TNFRSF10B. Briefly, recombinant human TRAIL were diluted serially in PBS, with 0.01%BSA (pH 7.4). Duplicate samples of 100uL were then transferred to TNFRSF10B-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-TRAIL pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50μ stop solution to the wells and read at 450nm immediately. The binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

activity of TRAIL and TNFRSF10B was shown, and this effect was in a dose dependent manner.

Molecular Mass	19.6kDa
Purity	> 95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	TNFSF10 tumor necrosis factor (ligand) superfamily, member 10 [Homo sapiens]
Official Symbol	TNFSF10
Synonyms	TNFSF10; tumor necrosis factor (ligand) superfamily, member 10; tumor necrosis factor ligand superfamily member 10; Apo 2L; CD253; TL2; TRAIL; Apo-2 ligand; TNF-related apoptosis inducing ligand TRAIL; tumor necrosis factor (ligand) family, member 10; chemokine tumor necrosis factor ligand superfamily member 10; tumor necrosis factor apoptosis-inducing ligand splice variant delta; APO2L; Apo-2L;
Gene ID	8743
mRNA Refseq	NM_001190942
Protein Refseq	NP_001177871

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


MIM 603598

UniProt ID P50591

Activity



SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA