

Active Recombinant Full Length Human PPM1A, His-tagged

Cat. No. PPM1A-66H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human PP2Cα was overexpressed in E. coli as a soluble His-tag fusion protein, and it was purified by conventional column chromatographic techniques, 46.6 kDa (418 aa).
Species	Human
Source	E.coli
Protein Length	1-382 a.a.
Description	Protein phosphatase 2C (PP2Cα) is a Mn ²⁺ - or Mg ²⁺ -dependent protein serine/threonine phosphatase that is essential for regulating cellular stress response in eukaryotes. The protein coding region of PP2Cα (amino acids 1-382) was cloned into an E. coli expression vector.
Sequences of amino acids	MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWILMGAF LDKPKMEKHN AQQQGNGLRY GLSSMQGWRV EMEDAHTAVI GLPSGLESWS FFAVYDGHAG SQVAKYCCEH LLDHITNNQD FKGSAGAPSV ENVKNGIRTG FLEIDEHMRV MSEKKHGADR SGSTAVGVLI SPQHTYFINC GDSRGLLCRN RKVHFFTQDH KPSNPLEKER IQNAGGSVMI QRVNGSLAVS RALGDFDYKC VHKGKPTQL VSPEPEVHDI ERSEDDQFI ILACDGIWDV MGNEELCDFV RSRLEVTDLL EKVCNEVVDV CLYKGSRDNM SVILICFPNA PKVSPEAVKK EAELDKYLEC RVEEIIKKQG EGVPLVHVM RTLASENIPS LPPGGELASK RNVIEAVYNR LNPYKNDDTD STSTDDMW

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Formulation	Liquid. In 10 mM Tris-HCl, pH7.5, 50 mM NaCl, 2 mM DTT, 1 mM MnCl ₂ , 20%glycerol.
Molecular Weight	46.6 kDa (418 aa), confirmed by MALDI-TOF
Purity	> 95% by SDS - PAGE
Concentration	1 mg/ml (determined by Bradford assay)
Endotoxin Level	< 1.0 EU per 1 µg of protein (determined by LAL method)
Specific Activity	>8,000 units/mg of PP2C alpha
Unit Difinition	One unit will hydrolyze 1nanomole of p-nitrophenyl phosphatate(pNPP) per minute at pH 7.4 at 37°C using 10mM of substrate
Storage	Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

GENE INFORMATION

Gene Name	PPM1A protein phosphatase 1A (formerly 2C), magnesium-dependent, alpha isoform [Homo sapiens]
Synonyms	Protein phosphatase 1A; EC 3.1.3.16, Protein phosphatase 2C isoform alpha; PP2C-alpha; IA; PPM1A; PP2CA; MGC9201; PPP2CA; PP2Ac; RP-C; protein phosphatase 2 (formerly 2A), catalytic subunit, alpha isoform; protein phosphatase 2, catalytic subunit, alpha isoform; PP2A-alpha; replication protein C; serine/threonine-protein phosphatase 2A catalytic subunit alpha isoform; FLJ42306; PP2C-ALPHA; PP2C-alpha; protein ph; protein phosphatase1A (formerly 2C), magnesium-dependent, alpha isoform

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	5494
mRNA Refseq	NM_021003
Protein Refseq	NP_066283
MIM	606108
UniProt ID	P35813
Chromosome Location	14q23.1
Function	hydrolase activity; magnesium ion binding; manganese ion binding; protein C-terminus binding; protein serine/threonine phosphatase activity; signal transducer activity

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA