

Active Recombinant Human ERBB2 Protein, Fc-tagged

Cat. No. ERBB2-033H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human HER2/ErbB2 Fc Chimera (rhHER2-Fc) Protein (AAA75493, 1-652aa), was expressed in human HEK293 cells with C-terminal human IgG1 Fc tag.
Species	Human
Source	HEK293
ProteinLength	1-652 aa
Description	This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized. [provided by RefSeq, Jul 2008]
Bio-activity	Measured by its ability to block anti-ErbB2 mediated inhibition of SK-BR-3 Human breast carcinoma cell proliferation. The ED50 for this effect is typically 20-80 ng/mL in

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	the presence of 0.6 µg/mL goat anti-hErbB2.
Molecular Mass	rhHER2-Fc was fused with Fc region of human IgG1 at C-terminal and has a calculated MW of 97.7 kDa. The predicted N-terminal is Thr 23. As a result of glycosylation, DTT-reduced protein migrates as 130-140 kDa polypeptide in SDS-PAGE.
Endotoxin	Less than 1.0 EU per 1 µg of the rhHER2-Fc by the LAL method.
Purity	>95% as determined by SDS-PAGE. All lots are greater than 95% pure.
Notes	Research Use Only. Not for use in diagnostic or therapeutic procedures.
Storage	Store at -20 centigrade in lyophilized state after receipt. For long term storage, upon reconstitution rhHER2-Fc should be aliquot and store at -20 centigrade or -8 centigrade. Avoid repeated freeze-thaw cycles.
Storage Buffer	Bulk protein in a 0.22 M filtered solution of PBS, pH 7.4 and delivered as liquid formulation or lyophilized powder. Normally 5-8% trehalose and mannitol are added as protectants before lyophilization.
GENE INFORMATION	
Gene Name	ERBB2 v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian) [Homo sapiens]
Official Symbol	ERBB2
Synonyms	ERBB2; v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian); NGL, v erb b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

oncogene homolog); receptor tyrosine-protein kinase erbB-2; CD340; HER 2; HER2; NEU; herstatin; p185erbB2; proto-oncogene Neu; c-erb B2/neu protein; proto-oncogene c-ErbB-2; metastatic lymph node gene 19 protein; tyrosine kinase-type cell surface receptor HER2; neuroblastoma/glioblastoma derived oncogene homolog; v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived oncogene homolog); NGL; TKR1; HER-2; MLN 19; HER-2/neu;

Gene ID	2064
mRNA Refseq	NM_001005862
Protein Refseq	NP_001005862
MIM	164870
UniProt ID	P04626