

## Recombinant Human BCL7C Protein, GST-tagged

**Cat. No.** BCL7C-166H    **Lot. No.** (See product label)

### SPECIFICATION

|                         |                                                                                                                                                                                                                                                                                                                           |
|-------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product Overview</b> | Human BCL7C full-length ORF (AAH19071.1, 1 a.a. - 217 a.a.) recombinant protein with GST-tag at N-terminal.                                                                                                                                                                                                               |
| <b>Species</b>          | Human                                                                                                                                                                                                                                                                                                                     |
| <b>Source</b>           | Wheat Germ                                                                                                                                                                                                                                                                                                                |
| <b>Description</b>      | This gene is identified by the similarity of its product to the N-terminal region of BCL7A protein. The BCL7A protein is encoded by the gene known to be directly involved in a three-way gene translocation in a Burkitt lymphoma cell line. The function of this gene has not yet been determined. [provided by RefSeq] |
| <b>Form</b>             | 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.                                                                                                                                                                                                                                                  |
| <b>Molecular Mass</b>   | 50.27 kDa                                                                                                                                                                                                                                                                                                                 |
| <b>AA Sequence</b>      | MAGRTVRAETRSRAKDDIKKVMATIEKVRRWEKRWVTVGDTSLRIFKWVPVDPQE<br>EERRRAGGGAERSRGRERRRGRGASPRGGGPLILLDLNDENSNQSFHSEGLQKGT<br>EPSPGGTPQSPRPVSPAGPPEGVPPEEAQPPRLGQERDPGGITAGSTDEPPMLTKE<br>EPVPELLEAEAPEAYPVFEPVPPVPEAAQGDTEDESEGAPPLKRICPNAPDP                                                                                    |
| <b>Notes</b>            | Best use within three months from the date of receipt of this protein.                                                                                                                                                                                                                                                    |
| <b>Storage</b>          | Store at -80 centigrade. Aliquot to avoid repeated freezing and thawing.                                                                                                                                                                                                                                                  |

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Storage Buffer** 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

## GENE INFORMATION

|                        |                                                                               |
|------------------------|-------------------------------------------------------------------------------|
| <b>Gene Name</b>       | BCL7C B-cell CLL/lymphoma 7C [ Homo sapiens ]                                 |
| <b>Official Symbol</b> | BCL7C                                                                         |
| <b>Synonyms</b>        | BCL7C; B-cell CLL/lymphoma 7C; B-cell CLL/lymphoma 7 protein family member C; |
| <b>Gene ID</b>         | 9274                                                                          |
| <b>mRNA Refseq</b>     | NM_004765                                                                     |
| <b>Protein Refseq</b>  | NP_004756                                                                     |
| <b>MIM</b>             | 605847                                                                        |
| <b>UniProt ID</b>      | Q8WUZ0                                                                        |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA