

Recombinant Bovine Fibroblast Growth Factor 2 (basic)

Cat. No. FGF2-1182B Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant bovine FGF2 is a single, non-glycosylated polypeptide chain containing 172 amino acids.
Species	Bovine
Source	E.coli
Description	FGF2 is a single-chain polypeptide growth factor that plays a significant role in the process of wound healing and is a potent inducer of angiogenesis. Several different forms of the human protein exist ranging from 18-24 kDa in size due to the use of alternative start sites within the FGF-2 gene. It has a 55 percent amino acid residue identity to FGF-1 and has potent heparin-binding activity.
Form	Lyophilized from a sterile filtered solution in PBS, pH 7.4 + 4% mannitol.
Molecular Weight	18.5 kDa
Purity	> 97% by SDS-PAGE and HPLC analyses
Endotoxin Level	< 0.1 ng/g of FGF2
Biological Activity	Calculated by the dose dependent proliferation of BALB/c 3T3 cells, the ED50 is <1.0 ng/ml.
Specific Activity	1 x 10 ⁶ Units/mg

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Sequences	MTMITNSSSVPGDPLESMAA GSITTLPALP EDGGSGAFPP GHFKDPKRLY CKNGGFFLRI HPDGRVDGVR EKSDPHIKLQLQAEERGVS IKGVCANRYL AMKEDGRLLA SKCVTDECFF FERLESNNYN TYRSRKYSSW YVALKRTGQYKLGPKTGPGQ KAILFLPMSA KS
Reconstitution	Centrifuge vial prior to opening. Add sterile distilled water or aqueous buffer to a concentration of 0.1-1.0 mg/mL. Further dilutions should be made in appropriate buffered solutions.
Storage	Stable at 2-8°C, but best kept desiccated -20°C. Upon reconstitution, stable for up to 1 week at 2-8°C. For longer term, store in working aliquots below -20°C. Avoid repeated freeze/thaw cycles.
Official Symbol	FGF2

GENE INFORMATION

Gene Name	FGF2 fibroblast growth factor2 (basic) [Bos taurus]
Synonyms	FGF2; fibroblast growth factor 2 (basic); heparin-binding growth factor 2; BFGF; HBGF-2; basic fibroblast growth factor
Gene ID	281161
mRNA Refseq	NM_174056
Protein Refseq	NP_776481
UniProt ID	P03969
Chromosome	17

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	Downstream signaling of activated FGFR; FGFR ligand binding and activation; FGFR1 ligand binding and activation; FGFR1b ligand binding and activation; FGFR1c ligand binding and activation; FGFR2 ligand binding and activation; FGFR2c ligand binding and activation; FGFR3 ligand binding and activation; FGFR4 ligand binding and activation; FRS2-mediated cascade; IRS-mediated signalling; Insulin receptor signalling cascade; MAPK signaling pathway; Melanoma
Function	growth factor activity; heparin binding; voltage-gated calcium channel activity
PDB rendering based on 1bas.	