

Native Human Coagulation Factor II (thrombin)

Cat. No. F2-5285H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Native Human F2 was expressed in human plasma.
Species	Human
Source	Human Plasma
Description	F2 is proteolytically cleaved to form thrombin in the first step of the coagulation cascade which ultimately results in the stemming of blood loss. F2 also plays a role in maintaining vascular integrity during development and postnatal life. Mutations in F2 lead to various forms of thrombosis and dysprothrombinemia.
Form	Liquid in 0.1 M NaH ₂ PO ₄ , pH 6.5.
Concentration	5.0 mg/ml
Molecular Weight	34.3 kDa
Specific Activity	Determined using a fibrinogen clotting assay.
Storage	Store in working aliquots at -20°C. Avoid repeated freeze-thaw cycles.
Pathways	Angiopoietin receptor Tie2-mediated signaling; Blood Clotting Cascade; Cell surface interactions at the vascular wall; Class A/1 (Rhodopsin-like receptors); Common Pathway; Complement and Coagulation Cascades; Diabetes pathways; FOXA2 and FOXA3 transcription factor networks; Extrinsic Pathway; Formation of

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

FibrinClot (Clotting Cascade); Hemostasis; Intrinsic Pathway; Regulation of actincytoskeleton; Selenium Pathway; Signal Transduction

GENE INFORMATION

Gene Name F2 coagulation factor II (thrombin) [Homosapiens]

Official Symbol F2

Synonyms F2; coagulation factor II (thrombin); PT; prothrombin; serineprotease; OTTHUMP00000197117; OTTHUMP00000197118; OTTHUMP00000233661; prothrombin B-chain; EC 3.4.21.5; EC 3.4.21; Prothrombin; Coagulation factorII; Activation peptide fragment 1; Activation peptide fragment 2; Thrombinlight chain; Thrombin heavy chain

Gene ID 2147

mRNA Refseq NM_000506

Protein Refseq NP_000497

MIM 176930

UniProt ID P00734

Chromosome Location 11p11


Function peptidaseactivity; protein binding; serine-type endopeptidase activity; heparin binding; calcium ion binding; growth factor activity; receptor binding; contributes toreceptor binging; thrombospondin receptor activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

The structure of
human thrombin in
complex with the
inhibitor hirudin.



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA