

Recombinant Human CD99 protein, His & T7-tagged

Cat. No. CD99-6941H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CD99 aa. (Ala31~Lys185 (Accession # P14209)) fused with N-terminal His & T7 tag was produced in E. coli cells.
Species	Human
Source	E.coli
ProteinLength	Ala31~Lys185
Description	The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte migration, T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. There is a related pseudogene located immediately adjacent to this locus.
Form	Freeze-dried powder
Molecular Mass	Predicted Molecular Mass: 19.6kDa
Endotoxin	<1.0EU per 1ug (determined by the LAL method)
Purity	>90%

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Characteristic	The isoelectric point is 4.7.
Applications	SDS-PAGE; WB; ELISA; IP
Stability	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Storage	Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.
Storage buffer	Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.
Reconstitution	Reconstitute in sterile PBS, pH7.2-pH7.4.

GENE INFORMATION

Gene Name	CD99 CD99 molecule (Xg blood group) [Homo sapiens (human)]
Official Symbol	CD99
Synonyms	CD99; CD99 molecule (Xg blood group); MIC2; HBA71; MIC2X; MIC2Y; MSK5X; CD99 antigen; E2 antigen; MIC2 (monoclonal antibody 12E7); T-cell surface glycoprotein E2; antigen identified by monoclonal 12E7, Y homolog; antigen identified by monoclonal antibodies 12E7, F21 and O13; cell surface antigen 12E7; cell surface antigen HBA-71; cell surface antigen O13; surface antigen MIC2
Gene ID	4267
mRNA Refseq	NM_001122898.2

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_001116370.1

UniProt ID P14209

SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA