

Transglutaminase Inhibitor Screening Assay Kit

Cat. No. Kit-2277 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Transglutaminases (TGs) are enzymes that catalyze post-translational modifications of glutamine residues in proteins, via the formation of isopeptide bonds, esterification and deamination. The reaction produces insoluble macromolecular complexes and protein aggregates. Within the TG family of enzyme, tissue-specific Transglutaminase-2 (TG2) expression is proven to be a clinical target for multiple types of cancer and neurodegenerative diseases, including Alzheimer's and Huntington's diseases. Inhibition of TG2, with a Cystamine-derived drug candidate (currently in phase-II clinical trial for Huntington's disease), has shown to have neuroprotective effects in models of neurodegeneration. Further, TG2 interacts with retinoblastoma protein (Rb), GTPase, and protein kinases through serine/threonine phosphorylation in cancer. Therefore, TG2 is a clinical validated candidate in these pathological diseases, and is a highly relevant therapeutic biomarker.

Transglutaminase Inhibitor Screening Assay Kit utilizes the catalytic activity of TG2, in a deamidation reaction between donor and acceptor substrates, to generate a hydroxamate product. The stop solution reacts with the hydroxamate product, forming a purple complex that can be measured at 525 nm. In the presence of TG2 inhibitor, the enzyme loses its ability to catalyze the deamidation reaction, leading to a decrease in color development and a lower absorbance OD reading. This assay kit is simple and can be used to identify and characterize TG2 inhibitors in a high-throughput format.

Applications	Screening/studying/characterizing Transglutaminase inhibitors.
---------------------	--

Storage	-20°C
----------------	-------

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Shipping	Gel Pack
Size	100 assays
Kit Components	TG Assay Buffer; 1 M DTT; Donor Substrate (lyophilized); Acceptor Substrate (lyophilized); Stop Solution; Transglutaminase-2 Enzyme (lyophilized); Inhibitor Cystamine (lyophilized)
Detection method	Absorbance (525 nm)
Features & Benefits	Simple method to screen Transglutaminase inhibitors; High-throughput Compatible; Includes Cystamine, an Inhibitor Control

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA