

HDAC2 Immunoprecipitation (IP) & Activity Assay Kit

Cat. No. Kit-2284 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Histone deacetylases (HDACs) play a central role in controlling cell cycle regulation, cell differentiation, and tissue development. These proteins have crucial roles in development and physiology, especially in the heart and nervous system. It is also deeply involved in cellular proliferation, cell cycle and apoptosis. The HDAC2 IP & Activity Assay Kit provides an antibody-based method to specifically immunoprecipitate the HDAC2 complex from cells & tissues and to measure HDAC2 activity fluorometrically. HDAC2 is immunoprecipitated from cell or nuclear extract(s) using HDAC2 antibody followed by capturing the complex on protein-A/G beads. The immunoprecipitated complex is incubated with the HDAC substrate. Only the deacetylated substrate is cleaved by the Developer to produce a fluorophore, which can be easily analyzed using a fluorescence plate reader.
Applications	Immunoprecipitation of HDAC2 complex from cell, nuclear, or tissue extract(s) Measurement of HDAC2 activity of immunoprecipitated complex/purified enzyme Screening for activators or inhibitors of HDAC2
Storage	-20°C
Shipping	Gel Pack
Size	25 assays
Kit Components	HDAC Assay Buffer; Cell Lysis Buffer; HDAC Substrate; Developer; AFC (7-amino-4-trifluoromethyl coumarin) Standard (1 mM); Rabbit HDAC2 Antibody (0.2 mg/ml);

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Rabbit IgG (Control Antibody) (0.5 mg/ml); Protein A/G Sepharose Beads; Positive Control (Jurkat Cell Lysate) (125 µg)

Detection method Fluorescence (Ex/Em=380/500 nm)

Features & Benefits Simple, fast and convenient.
Only kit on the market for both IP as well as measuring activity of HDAC2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA