

Active Recombinant Mouse Junction Adhesion Molecule 3, Fc-tagged

Cat. No. Jam3-5667M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Jam3 protein was expressed in Murine myeloma cell line with the Fc part of Human IgG1 Pro100-Lys330(C-terminus) and Mouse Jam3 Val32-Asn241(N-terminus).
Species	Mouse
Source	Mammalian Cells
Protein Length	32-241 a.a.
Description	The family of junctional adhesion molecules (JAM), comprised of at least three members, are type I transmembrane receptors belonging to the immunoglobulin(Ig) superfamily. These proteins are localized in the tight junctions between endothelial cells or epithelial cells. Some family members are also found on blood leukocytes and platelets. Mouse Jam3 cDNA predicts a 310 amino acid (aa) residue precursor protein with a putative 31aa signal peptide, a 210 aa extracellular region containing two Ig domains, an 18 aa transmembrane domain and a 51 aa cytoplasmic domain containing a PDZ-binding motif and a PKC phosphorylation site.
Form	Lyophilized from a 0.2 µm filtered solution in Tris-Citrate and NaCl.
N-terminal Sequence	Val32
Molecular Weight	50 kDa (monomer)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Activity	Measured by its ability to inhibit the adhesion of J45.01 human acute T lymphoblastic leukemia cells on immobilized JAM-2 (VE-JAM)/Fc Chimera. The ED50 for this effect is typically 0.1-0.5 µg/mL in the presence of 1x10 ⁵ cells/well. Each well was coated with 100 µL of a 0.2 µg/mL solution of rhJAM-2(VE-JAM)/Fc Chimera.
Endotoxin Level	<1.0 EU per 1 µg of the protein by the LAL method.
SDS-PAGE	64 kDa, reducing conditions.
Purity	>95%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Reconstitution	100 µg/mL in sterile PBS
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 1 month, 2 to 8°C under sterile conditions after reconstitution. 3 months, -20 to -70°C under sterile conditions after reconstitution.

GENE INFORMATION

Gene Name	Jam3 junction adhesion molecule 3 [<i>Mus musculus</i>]
Official Symbol	Jam3
Synonyms	Jam3; junction adhesion molecule 3; JAM-2; junction cell adhesion molecule 3; junction cell adhesion molecule C; junctional adhesion molecule 3; junctional adhesion molecule C; 1110002N23Rik, C85515, JAM-3, JAM-C
Gene ID	83964
mRNA Refseq	NM_023277

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_075766
UniProt ID	Q9D8B7
Chromosome Location	9
Pathway	Cell adhesion molecules (CAMs); Cell surface interactions at the vascular wall; Hemostasis; Integrin cell surface interactions; Leukocyte transendothelial migration; Signal Transduction; Tight junction
Function	integrin binding; protein binding; protein heterodimerization activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA