

Active Recombinant Human Interleukin 4 Induced 1, His-tagged

Cat. No. IL4I1-654H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL4I1 protein was expressed in Chinese Hamster Ovary cell line. Gln22-His567, with a C-terminal 6-His tag.
Species	Human
Source	CHO
Protein Length	22-567 a.a.
Description	IL4I1 is encoded by a B-cell IL-4-inducible gene, FIG1, and is highly expressed in primary metastatic B-cell lymphomas. It belongs to the flavin monoamine oxidase family, FIG1 subfamily. Enzymological characterization reveals that IL4I1 has L-amino acid oxidase activity with preference toward aromatic amino acids. Studies have shown that hIL-4I1 inhibited the proliferation of CD3-stimulated T lymphocytes with a similar effect on CD4(+) and CD8(+) T cells. Its inhibitory effect was dependent on enzymatic activity and H ₂ O ₂ production. Its restricted expression to lymphoid tissues indicates that it may play an important function in the immune system.
Form	Supplied as a 0.2 umfiltered solution in Tris, NaCl and Glycerol.
N-terminal Sequence	Gln22 predicted.
Molecular Weight	61 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Activity	Measured by its ability to oxidize phenylalanine in a horseradish peroxidase coupled assay. The specific activity is > 150 pmol/min/μg, as measured under the described conditions.
Endotoxin Level	<1.0 EU per 1 μg of the protein by the LAL method.
SDS-PAGE	70-75 kDa, reducing conditions.
Purity	>95%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 6 months from date of receipt, -20 to -70°C as supplied. 3 months, -20 to -70°C under sterile conditions after opening.

GENE INFORMATION

Gene Name	IL4I1 interleukin 4 induced 1 [Homo sapiens]
Official Symbol	IL4I1
Synonyms	IL4I1; interleukin 4 induced 1; LAO; FIG1; LAAO; L-amino-acid oxidase; Fig-1 protein; IL4-induced protein 1; interleukin four induced 1; EC1.4.3.2
Gene ID	259307
mRNA Refseq	NM_172374
Protein Refseq	NP_758962
MIM	609742

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q96RQ9
Chromosome Location	19q13.3-q13.4
Pathway	Alanine, aspartate and glutamate metabolism;Cysteine and methionine metabolism; Metabolic pathways; Phenylalaninemetabolism; Phenylalanine; Tryptophan metabolism; Tyrosine metabolism; Valine,leucine and isoleucine degradation
Function	L-amino-acid oxidaseactivity; oxidoreductase activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA