

Active Recombinant Rat Chemokine (C-X-C motif) Ligand 5

Cat. No. Cxcl5-445R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Rat Cxcl5 protein was expressed in E. coli. Ala38-Arg125& Ala38-Lys117.
Species	Rat
Source	E.coli
Protein Length	38-125 a.a.
Description	Cxcl5 is a chemokine originally cloned from LPS stimulated mouse fibroblasts. Rat Cxcl5 shares approximately 74% amino acid (aa) sequence identity with mouse Cxcl5 and is likely the rat orthologue to mouse Cxcl5. Rat Cxcl5 cDNA encodes a 130 aa residue precursor with a predicted 37 aa residue signal peptide and a 93 aa residue mature protein. Among human CXC chemokines, rat Cxcl5 is most closely related to human GCP2 and ENA78. Rat Cxcl5 also differs from these two human proteins by having an extended carboxy terminus.
Form	Lyophilized from a 0.2 µm filtered solution in Acetonitrile and TFA with BSA as a carrier protein.
N-terminal Sequence	Ala38
Molecular Weight	10.1 kDa
Activity	Measured by its ability to chemoattract human CXCR2 transfected BaF3 mouse proB

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	cells. The ED50 for this effect is typically 0.03-0.15 µg/mL.
Endotoxin Level	<1.0 EU per 1 µg of the protein by the LAL method.
Purity	>97%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Reconstitution	Reconstitute at 100 µg/mL in sterile PBS containing at least 0.1% human or bovine serum albumin.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, 20 to 70°C as supplied. 1 month, 2 to 8°C under sterile conditions after reconstitution. 3 months, 20 to 70°C under sterile conditions after reconstitution.

GENE INFORMATION

Gene Name	Cxcl5 chemokine (C-X-C motif) ligand 5 [<i>Rattus norvegicus</i>]
Official Symbol	Cxcl5
Synonyms	Cxcl5; chemokine (C-X-C motif) ligand 5; Cxcl6; C-X-C motif chemokine 5; cytokine LIX; CXC chemokine LIX; small-inducible cytokine B5; chemokine (C-X-C motif) ligand 6 (granulocyte chemotactic protein 2)
Gene ID	60665
mRNA Refseq	NM_022214
Protein Refseq	NP_071550
UniProt ID	P97885

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	14
Pathway	Chemokine receptorsbind chemokines; Chemokine signaling pathway; Class A/1 (Rhodopsin-likereceptors); G alpha (i) signalling events; GPCR downstream signaling; GPCRligand binding; Peptide ligand-binding receptors; Pertussis; Rheumatoidarthritis; Signal Transduction; Signaling by GPCR
Function	chemokine activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA