

Active Recombinant Mouse Chemokine (C-C motif) Ligand 17

Cat. No. Ccl17-463M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Ccl17protein was expressed in E. coli. Ala24-Pro93.
Species	Mouse
Source	E.coli
ProteinLength	24-93 a.a.
Description	Human thymus and activationregulatedchemokine (TARC) also known as Ccl17, is a novel CC chemokine identified using a signal sequence trap method. Mouse Ccl17 was discovered as a dendritic cell(DC) specific gene by differentiation RNA display. Mouse Ccl17 cDNA encodes a highly basic 93 amino acid (aa) residue precursor protein with a 23 aa residueputative signal peptide that is cleaved to generate the 70 aa residue mature secretproteins.
Form	Lyophilized from a 0.2µm filtered solution in Acetonitrile and TFA with BSA as a carrier protein.
N-terminalSequence	Ala24
Molecular Weight	7.9 kDa
Activity	Measuredby its ability to chemoattract CEMNKR human Tlymphoblastoid cells. Howell, D.N.et al. (1985) J. Immunol. 134:971. The ED50 for this effect is typically2 10 ng/mL. Measured by its ability to chemoattract human CCR4 transfected

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

BaF3mouse proB cells. The ED50 for this effect is typically 2-8 ng/mL.

Endotoxin Level <1.0 EU per 1 µg of the protein by the LAL method.

Purity >97%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Reconstitution Reconstitute at 25 µg/mL in sterile PBS that containing at least 0.1% human or bovine serum albumin.

Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 1 month, 2 to 8°C under sterile conditions after reconstitution. 3 months, -20 to -70°C under sterile conditions after reconstitution.

GENE INFORMATION

Gene Name Ccl17 chemokine(C-C motif) ligand 17 [*Mus musculus*]

Official Symbol Ccl17

Synonyms Ccl17; chemokine(C-C motif) ligand 17; Tarc; Abcd-2; Scya17; Scya17l; small inducible cytokine subfamily A17; CC chemokine ABCD-2; Thymus and activation-regulated chemokine; mCG_12286

Gene ID 20295

mRNA Refseq NM_011332

Protein Refseq NP_035462

UniProt ID Q9WUZ6


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	8 C5; 8 45.0 cM
Pathway	Chemokine signaling pathway;Cytokine-cytokine receptor interaction
Function	CCR4 chemokine receptorbinding; chemokine activity; cytokine activity

PDB renderingbased
on 1nr2.



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA