

Recombinant Human CST9L Protein, MYC/DDK-tagged

Cat. No. CST9L-3351H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human CST9L protein, fused to MYC/DDK-tagged at C-terminus, was expressed in HEK293.
Species	Human
Source	HEK293
Description	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a protein similar to mouse cystatin 9. Based on its testis-specific expression, it is likely to have a role in tissue reorganization during early testis development.
Form	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.
Molecular Mass	14.2 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration	>50 ug/mL as determined by microplate BCA method

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GENE INFORMATION

Gene Name	CST9L cystatin 9-like [Homo sapiens]
Official Symbol	CST9L
Synonyms	bA218C14.1; CTES7B
Gene ID	128821
mRNA Refseq	NM_080610
Protein Refseq	NP_542177
UniProt ID	Q9H4G1

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA