

Recombinant Human CST2 Protein, MYC/DDK-tagged

Cat. No. CST2-3357H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CST2 protein, fused to MYC/DDK-tagged at C-terminus, was expressed in HEK293.
Species	Human
Source	HEK293
Description	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a secreted thiol protease inhibitor found at high levels in saliva, tears and seminal plasma
Form	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.
Molecular Mass	14.3 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration	>50 ug/mL as determined by microplate BCA method

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GENE INFORMATION

Gene Name	CST2 cystatin SA [Homo sapiens]
Official Symbol	CST2
Synonyms	MGC71924
Gene ID	1470
mRNA Refseq	NM_001322
Protein Refseq	NP_001313
MIM	123856
UniProt ID	P09228

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA