

Active Recombinant Human FGF4 Protein

Cat. No. FGF4-4113H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human FGF4 (P08620) recombinant protein expressed in Escherichia coli.
Species	Human
Source	E.coli
Description	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This gene was identified by its oncogenic transforming activity. This gene and FGF3, another oncogenic growth factor, are located closely on chromosome 11. Co-amplification of both genes was found in various kinds of human tumors. Studies on the mouse homolog suggested a function in bone morphogenesis and limb development through the sonic hedgehog (SHH) signaling pathway. [provided by RefSeq]
Form	Lyophilized
Bio-activity	The activity is determined by the ability to induce the proliferation of mouse NR6R-3T3 fibroblasts. The expected ED50 for this effect is less than 0.1 ng/mL.
Molecular Mass	19 kDa
AA Sequence	MAPTAPNGTLEAELERRWESLVALSLARLPVAAQPKEAAVQSGAGDYLLGIKRLRRL YCNVGIGFHLQALPDGRIGGAHADTRDSLLELSPVERGVVSIFGVASRFFVAMSSKG

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	KLYGSPFFTDLECTFKEILLPNNYNAYESYKYPGMFIALSKNGKTKKGNRVSPMKVT HFLPRL
Endotoxin	< 0.1 EU/μg
Applications	Functional Study SDS-PAGE
Storage	Store at -20 centigrade on dry atmosphere. After reconstitution with sterilized water, store at -20 centigrade or lower. Aliquot to avoid repeated freezing and thawing.
Storage Buffer	Lyophilized from 0.5X PBS, pH 7.5

GENE INFORMATION

Gene Name	FGF4 fibroblast growth factor 4 [Homo sapiens]
Official Symbol	FGF4
Synonyms	FGF4; fibroblast growth factor 4; heparin secretory transforming protein 1, HSTF1; HBGF 4; HST; HST 1; human stomach cancer; transforming factor from FGF related oncogene; K FGF; kaposi sarcoma oncogene; KFGF; transforming protein KS3; FGF-4; HSTF-1; oncogene HST; heparin-binding growth factor 4; heparin secretory transforming protein 1; heparin secretory-transforming protein 1; fibroblast growth factor 4 splice isoform; human stomach cancer, transforming factor from FGF-related oncogene; HST-1; HSTF1; K-FGF; HBGF-4;
Gene ID	2249
mRNA Refseq	NM_002007

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_001998

MIM 164980

UniProt ID P08620

Result of activity analysis



3T3 cells were cultured with 0 to 1 µg/mL human FGF4. Cell proliferation was measured after 42 hours and the linear portion of the curve was used to calculate the ED50.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA