

Recombinant Human Histone Deacetylase 6, FLAG-tagged

Cat. No. HDAC6-398H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinanthuman HDAC6 was expressed in E. coli as the full length protein with anN-terminal FLAG tag. The molecular weight of the protein is 132 kDa.
Species	Human
Source	E.coli
Description	HDAC6is a histone deacetylase containig two HDAC catalytic domains. HDAC6 is anenzyme that in humans is encoded by the HDAC6 gene. Histones play a critical rolein transcriptional regulation, cell cycle progression, and developmentalevents. It contains an internal duplication of two catalytic domains whichappear to function independently of each other.
Reconstitution	20 mM Tris-HCl (pH7.3), 300 mM KCl, 0.2 mM EDTA and 20%glycerol.
Contents	10g recombinant HDAC6 protein, 0.5 g/l.
Dilution of Protein	RecombinantHDAC6 is supplied in solution and should be kept as a solution in ordertomaintain full activity.
Application	RecombinantHDAC6 is suitable for use in histone deacetylase assays. Arecommendedstarting point for the HDAC assay is 20-100 ng of recombinant proteinperreaction.
Storage	Recombinantproteins in solution are temperature sensitive and should be stored at-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

80°C to prevent degradation. Avoid repeated freeze/thaw cycles and keep on ice when not in storage.

Full Length Full L.

GENE INFORMATION

Gene Name HDAC6 histone deacetylase 6 [Homo sapiens]

Official Symbol HDAC6

Synonyms HDAC6; histone deacetylase 6; HD6; JM21; FLJ16239; EC 3.5.1.98; KIAA0901; OTTHUMP00000032398; OTTHUMP00000197663

Gene ID 10013

mRNA Refseq NM_006044

Protein Refseq NP_006035

MIM 300272

UniProt ID Q9UBN7

Chromosome Location Xp11.23

Pathway Cell cycle; NOTCH1 Intracellular Domain Regulates Transcription; Signal Transduction; Signaling by NOTCH; Signaling by NOTCH1; Signaling events mediated by HDAC Class I; Signaling events mediated by HDAC Class II; TNF-alpha/NF-kB Signaling Pathway

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

Hsp90 protein binding; NAD-dependent histone deacetylase activity (H3-K14 specific); NAD-dependent histone deacetylase activity (H3-K9 specific); NAD-dependent histone deacetylase activity (H4-K16 specific); actin binding; alpha-tubulin binding; beta-catenin binding; beta-tubulin binding; dynein complex binding; enzyme binding; histone deacetylase activity; histone deacetylase activity (H3-K16 specific); histone deacetylase binding; hydrolase activity; metal ion binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA