

Recombinant Mouse TNFSF14 protein

Cat. No. TNFSF14-2020M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse TNFSF14 protein was expressed in Escherichia coli.
Species	Mouse
Source	E.coli
ProteinLength	168
Description	The protein encoded by this gene is a member of the tumor necrosis factor (TNF) ligand family. This protein is a ligand for TNFRSF14, which is a member of the tumor necrosis factor receptor superfamily, and which is also known as a herpesvirus entry mediator (HVEM). This protein may function as a costimulatory factor for the activation of lymphoid cells and as a deterrent to infection by herpesvirus. This protein has been shown to stimulate the proliferation of T cells, and trigger apoptosis of various tumor cells. This protein is also reported to prevent tumor necrosis factor alpha mediated apoptosis in primary hepatocyte. Two alternatively spliced transcript variant encoding distinct isoforms have been reported.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cytotoxicity assay using human HT-29 cells is less than 2 µg/ml, corresponding to a specific activity of > 500 IU/mg in the presence of murine anti-polyHistidine monoclonal antibody and rHuIFN-γ.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 18.4 kDa, a single non-glycosylated polypeptide chain containing 168 amino acids.
AA Sequence	DGGKGSWEKLIQDQRSHQANPAAHLTGANASLIGIGGPLLWETRLGLAFLRGLTYH DGALVTMEPGYYYVYSKVQLSGVGCPQGLANGLPITHGLYKRTSRYPKELELLVSR RSPCGRANSSRVWWDSSFLGGVVHLEAGEEVVVRVPGNRLVRPRDGTRSYFGAF MV
Endotoxin	Less than 1 EU/g of rMuLIGHT/TNFSF14 as determined by LAL method.
Purity	>96% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 100 mM HAc to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	TNFSF14
Official Symbol	TNFSF14
Synonyms	TNFSF14; tumor necrosis factor (ligand) superfamily, member 14; tumor necrosis factor ligand superfamily member 14; LTg; HVEM-L; LIGHT; Ly113; HVEM-L;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Gene ID 50930

mRNA Refseq NM_019418

Protein Refseq NP_062291

UniProt ID Q9QYH9

**SDS-PAGE of
TNFSF14-2020M**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA