

Recombinant Human BAG3 Protein, MYC/DDK-tagged

Cat. No. BAG3-2138H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human BAG3 protein, fused to MYC/DDK-tagged at C-terminus, was expressed in HEK293
Species	Human
Source	HEK293
Description	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq, Jul 2008].
Form	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.
Molecular Mass	61.4 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration	>50 ug/mL as determined by microplate BCA method

GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	BAG3 BCL2-associated athanogene 3 [Homo sapiens]
Official Symbol	BAG3
Synonyms	BAG-3; BIS; CAIR-1; MFM6
Gene ID	9531
mRNA Refseq	NM_004281
Protein Refseq	NP_004272
MIM	603883
UniProt ID	O95817

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA