

Mouse Anti-Human Clusterin Monoclonal Antibody

Cat. No. CAB11516MH **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse Monoclonal Antibody to Human Clusterin
Species	Human
Source	Mouse
Antigen Description	Clusterin, also known as complement-associated protein SP-40, Complement cytolysis inhibitor, Apolipoprotein J, Testosterone-repressed prostate message 2, Aging-associated gene 4 protein, CLU and APOJ, is a secreted protein which belongs to the clusterin family. Clusterin is an enigmatic glycoprotein with a nearly ubiquitous tissue distribution and an apparent involvement in biological processes ranging from mammary gland involution to neurodegeneration in Alzheimers disease. Its major form, a heterodimer, is secreted and present in physiological fluids, but truncated forms targeted to the nucleus have also been identified. CLU / Clusterin is a widely distributed glycoprotein with a wide range of biologic properties. A prominent and defining feature of clusterin is its marked induction in such disease states as glomerulonephritis, cystic renal disease, renal tubular injury, neurodegenerative conditions, atherosclerosis, and myocardial infarction. Upregulation of clusterin mRNA and protein levels detected in diverse disease states and in in vitro systems have led to suggestions that it functions in membrane lipid recycling, in apoptotic cell death, and as a stress-induced secreted chaperone protein, amongst others.
Specificity	Human Clusterin / Apolipoprotein J / Apo-J. No cross-reactivity in ELISA with Mouse CLU / Clusterin / Apolipoprotein J / Apo-J; Human cell lysate (293 cell line)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Immunogen	Recombinant Human Clusterin / Apolipoprotein J / Apo-J Protein
Isotype	Mouse IgG1
Clone	10F3F3
Applications	WB; ELISA
Dilution	Western blot: This antibody can be used at 1-2 µg/mL with the appropriate secondary reagents to detect Human CLU in WB. Using a DAB detection system, the detection limit for Human CLU is approximately 20 ng/lane under non-reducing conditions and reducing conditions.ELISA: This antibody can be used at 0.5-1 µg with the appropriate secondary reagents to detect Human CLU. The detection limit for Human CLU is approximately 0.16 ng/well.
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, human cell-derived, recombinant Human Clusterin / Apolipoprotein J / Apo-J. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Format	0.2 µm filtered solution in PBS with 5% trehalose
Storage	This antibody can be stored at 2-8 °C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 to -70°C. Preservative-Free.Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

GENE INFORMATION

Gene Name	CLU clusterin [Homo sapiens]
------------------	--------------------------------

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	CLU
Synonyms	CLU; clusterin; APOJ, CLI, clusterin (complement lysis inhibitor, SP 40,40, sulfated glycoprotein 2, testosterone repressed prostate message 2, apolipoprotein J); apolipoprotein J; complement lysis inhibitor; KUB1; SGP 2; SP 40; sulfated glycoprotein 2; testosterone repressed prostate message 2; TRPM 2; ku70-binding protein 1; aging-associated protein 4; complement cytolysis inhibitor; complement-associated protein SP-40,40; testosterone-repressed prostate message 2; CLI; AAG4; APOJ; SGP2; APO-J; SGP-2; SP-40; TRPM2; TRPM-2; NA1/NA2; MGC24903;
Gene ID	1191
mRNA Refseq	NM_001831
Protein Refseq	NP_001822
MIM	185430
UniProt ID	P10909
Chromosome Location	8p21-p12
Pathway	Hemostasis, organism-specific biosystem; Platelet activation, signaling and aggregation, organism-specific biosystem; Platelet degranulation, organism-specific biosystem; Response to elevated platelet cytosolic Ca ²⁺ , organism-specific biosystem; Validated targets of C-MYC transcriptional repression, organism-specific biosystem;
Function	NOT ATPase activity; misfolded protein binding; misfolded protein binding; protein

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

binding; ubiquitin protein ligase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA