

Rabbit Anti-Mouse Cystatin C Monoclonal Antibody

Cat. No. CAB11520RM **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Rabbit Monoclonal Antibody to Mouse Cystatin C
Species	Mouse
Source	Rabbit
Antigen Description	Mouse Cystatin-C, also known as Cystatin-3, Neuroendocrine basic polypeptide, Post-gamma-globulin, and CST3, is a secreted type 2 cysteine proteinase inhibitors of the cystatin superfamily, and exerts actions on a variety of cysteine proteases including papain and Cathepsins B, H, K, L and S. As the most abundant extracellular inhibitor of cysteine proteases, Cystatin-C encoded by a housekeeping gene is found ubiquitously in all organs and biological fluids at high concentration levels. It has been shown that Cystatin-C binds amyloid-beta and reduces cerebral amyloid-beta deposition or inhibits its fibril formation in an APP-transgenic mouse model. Thus, a point mutation in this gene (Leu 68 to Gln) has been associated with a severe hereditary form of amyloid angiopathy. Studies have also investigated Cystatin-C as a marker of kidney function and thyroid dysfunction prior to creatinine, and could be a useful clinical tool to identify HIV-infected persons. In addition, Cystatin-C expression is up-regulated in malignance of certain tumor progression.
Specificity	Mouse Cystatin C / CST3. No cross-reactivity in ELISA with Human CST3; Human HPRG; Human KNG1; Human cell lysate (293 cell line);
Immunogen	Recombinant Mouse Cystatin C / CST3 Protein

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Isotype	Rabbit IgG
Clone	A042
Applications	WB; ELISA
Dilution	Western blot: This antibody can be used at 1-2 µg/mL with the appropriate secondary reagents to detect Mouse CST3 in WB. Using a DAB detection system, the detection limit for Mouse CST3 is approximately 0.5 ng/lane under non-reducing conditions and 0.25 ng/lane under reducing conditions. ELISA: This antibody can be used at 0.1-0.2 µg/mL with the appropriate secondary reagents to detect Mouse CST3. The detection limit for Mouse CST3 is approximately 0.00245 ng/well.
Preparation	This antibody was obtained from a rabbit immunized with purified, human cell-derived, recombinant Mouse Cystatin C / CST3
Format	0.2 µm filtered solution in PBS with 5% trehalose
Storage	This antibody can be stored at 2-8 °C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 to -70°C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

GENE INFORMATION

Gene Name	Cst3 cystatin C [<i>Mus musculus</i>]
Official Symbol	Cst3
Synonyms	CST3; cystatin C; cystatin-C; cystatin 3; cystatin-3; CysC;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	13010
mRNA Refseq	NM_009976
Protein Refseq	NP_034106
Pathway	Salivary secretion, organism-specific biosystem; Salivary secretion, conserved biosystem;
Function	beta-amyloid binding; cysteine-type endopeptidase inhibitor activity; endopeptidase inhibitor activity; peptidase inhibitor activity; protease binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA