

Rabbit Monoclonal Antibody to Human Serine Peptidase Inhibitor, Clade F, Member 1

Cat. No. CAB11602RH Lot. No. (See product label)

SPECIFICATION

Product Overview	Rabbit Monoclonal Antibody to Human Serine Peptidase Inhibitor, Clade F, Member 1
Species	Human
Source	Rabbit
Antigen Description	Pigment epithelium-derived factor, also known as PEDF, Serpin F1, and SERPINF1, is a multiple functional protein which has both anti-angiogenic activity and neurotrophic activity at the same time. PEDF is a secreted glycoprotein that belongs to the noninhibitory serpin. It has an alpha/beta core serine-protease inhibitor domain, three major beta-sheets, and ten alpha-helices. PEDF does not inhibit either serine or cysteine proteinases. PEDF exerts diverse physiological activities including anti-angiogenesis, anti-vasopermeability, anti-tumor, and neurotrophic activities. PEDF acts via multiple high affinity ligands and cell receptors. It has been described as a natural angiogenesis inhibitor with neurotrophic and immune-modulation properties. PEDF induces macrophages apoptosis and necrosis through the activation of peroxisome proliferator-activated receptor-gamma by which PEDF could modulate inflammatory reactions in septic shock. It balances angiogenesis in the eye and blocks tumor progression.
Specificity	Human PEDF / SerpinF1. No cross-reactivity in ELISA with Mouse PEDF / SerpinF1; Human cell lysate (293 cell line)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Immunogen	Recombinant Human PEDF / SerpinF1 Protein
Isotype	Rabbit IgG
Clone	A014
Applications	Western blot; ELISA
Dilution	Western blot: This antibody can be used at 1-2 µg/mL with the appropriate secondary reagents to detect Human SERPINF1 in WB. Using a DAB detection system, the detection limit for Human SERPINF1 is approximately 0.5 ng/lane under non-reducing conditions and 0.25 ng/lane under reducing conditions. Direct ELISA: This antibody can be used at 0.1-0.2 µg/mL with the appropriate secondary reagents to detect Human SERPINF1. The detection limit for Human SERPINF1 is approximately 0.00975 ng/well.
Preparation	This antibody was obtained from a rabbit immunized with purified, human cell-derived, recombinant Human PEDF / SerpinF1
Format	0.2 µm filtered solution in PBS with 5% trehalose
Storage	This antibody can be stored at 2-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 to -70 °C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

GENE INFORMATION

Gene Name SERPINF1 serpin peptidase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1 [Homo sapiens]

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	SERPINF1
Synonyms	SERPINF1; serpin peptidase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1; PEDF,serine (or cysteine) proteinase inhibitor, clade F (alpha 2 antiplasmin, pigment epithelium derived factor), member 1; pigment epithelium-derived factor; EPC 1; PIG35; pigment epithelium derived factor; proliferation inducing protein 35; serpin F1; cell proliferation-inducing gene 35 protein; serine (or cysteine) proteinase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1; OI6; OI12; PEDF; EPC-1;
Gene ID	5176
mRNA Refseq	NM_002615
Protein Refseq	NP_002606
MIM	172860
UniProt ID	P36955
Chromosome Location	17p13.3
Function	serine-type endopeptidase inhibitor activity;