

Recombinant Human UPF3B Protein, MYC/DDK-tagged

Cat. No. UPF3B-631H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human UPF3B protein, fused to MYC/DDK tag at C-terminus, was expressed in HEK293.
Species	Human
Source	HEK293
Description	This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. The encoded protein is one of two functional homologs to yeast Upf3p. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein binds to the mRNA and remains bound after nuclear export, acting as a nucleocytoplasmic shuttling protein. It forms with Y14 a complex that binds specifically 20 nt upstream of exon-exon junctions. This gene is located on the long arm of chromosome X. Two splice variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008].
Form	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Molecular Mass	56 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Concentration >50 ug/mL as determined by microplate BCA method

GENE INFORMATION

Gene Name	UPF3B, regulator of nonsense mediated mRNA decay[Homo sapiens]
Official Symbol	UPF3B
Synonyms	HUPF3B; MRX62; MRXS14; RENT3B; UPF3BP1; UPF3BP2; UPF3BP3; Upf3p-X; UPF3X
Gene ID	65109
mRNA Refseq	NM_080632.2
Protein Refseq	NP_542199.1
MIM	300298
UniProt ID	Q9BZI7

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA