

Recombinant Human Proteasome (prosome, macropain) Activator Subunit 3 (PA28 gamma; Ki), His-tagged

Cat. No. PSME3-6860H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human PSME3 protein, fused to His-tag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques. MW: 31.7 kDa.
Species	Human
Source	E.coli
ProteinLength	1-254aa
Description	PSME3 belongs to the PA28 family. The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. PSME3 activates the trypsin-like catalytic subunit of the proteasome but inhibits the chymotrypsin-like and postglutamyl-preferring subunits.
Form	Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 2mM DTT, 40% glycerol, 200mM NaCl.
Molecular Mass	31.7kDa (274aa), confirmed by MALDI-TOF.
AA Sequence	MGSSHHHHHH SSGLVPRGSH MASLLKVDQE VKLKVDSFRE RITSEAEDLV ANFFPKKLL LDSFLKEPIL NIHDLTQIHS DMNLPVPDPI LLTNSHDGLD

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GPTYKKRRLD ECEEAFQGTK VFVMPNGMLK SNQQLVDIIE KVKPEIRLLI
 EKCNTVKMWV QLLIPRIEDG NNFGVSIQEE TVAELRTVES EAASYLDQIS
 RYYITRAKLV SKIAKYPHVE DYRRTVTEID EKEYISLRLLI ISELRNQYVT
 LHDMLKNIE KIKRPRSSNA ETLV

Purity	>90% by SDS - PAGE
Applications	SDS-PAGE
Storage	Can be stored at 4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.
Concentration	0.5mg/ml (determined by Bradford assay)

GENE INFORMATION

Gene Name	PSME3 proteasome (prosome, macropain) activator subunit 3 (PA28 gamma; Ki) [Homo sapiens]
Official Symbol	PSME3
Synonyms	PSME3; proteasome (prosome, macropain) activator subunit 3 (PA28 gamma; Ki); proteasome activator complex subunit 3; Ki; PA28 gamma; PA28G; REG GAMMA; PA28gamma; Ki antigen; Ki nuclear autoantigen; proteasome activator 28-gamma; 11S regulator complex gamma subunit; 11S regulator complex subunit gamma; proteasome activator 28 subunit gamma; activator of multicatalytic protease subunit 3; REG-GAMMA; PA28-gamma;
Gene ID	10197
mRNA Refseq	NM_005789

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_005780
MIM	605129
UniProt ID	P61289
Chromosome Location	17q12-q21
Pathway	APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem; APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem;
Function	MDM2 binding; endopeptidase activator activity; p53 binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA