

Active Recombinant Human Colony Stimulating Factor 3 (Granulocyte)

Cat. No. CSF3-142H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Colony Stimulating Factor 3 (Granulocyte) encoding the human Colony Stimulating Factor 3 (Granulocyte) protein sequence (containing the signal peptide sequence, and the mature G-CSF sequence) was expressed in modified <i>human 293 cells</i> .
Species	Human
Source	HEK293
Description	Granulocyte colony stimulating factor (G-CSF) is the primary extracellular regulator of granulopoiesis and regulates the production of neutrophils by stimulating proliferation and survival of specific bone marrow precursor cells and their differentiation into granulocytes. Neutrophils play a critical role in the defence against bacterial and fungal infections. G-CSF is produced by monocytes, macrophages, neutrophils, fibroblasts and endothelial cells and is capable of increasing the absolute number of circulating neutrophils and enhancing their antimicrobial function. Unlike GM-CSF, the activity of G-CSF is not species specific. Additionally, G-CSF production is inducible by cytokines including TNF-alpha, IL-1, GM-CSF, IL-17 and IL-4.
Amino Acid Sequence	TPLGPASSLPQSFLKCLEQVRKIQGDGAALQEKLCATYKLCHPEELVLLGHSLGIP WAPLSSCPSQALQLAGCLSQLHSGFLYQGILLQALEGISPELGPTLDTLQLDVADFA TTIWQQMEELGMAPALQPTQGAMPAFASAFQRRAGGVLVASHLQSFLEVSYRVLR HLAQP.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	G-CSF migrates as a band between 15 and 20 kDa in SDS-PAGE. This compares with the predicted molecular mass of 18.7 kDa.
pI	G-CSF separates into a number of isoforms with a pI between 5.4 and 6.0 in 2D PAGE Due to post-translational modifications, in particular glycosylation.
Purity	>95%, as determined by SDS-PAGE and visualized by silver stain.
Formulation	When reconstituted in 0.5 ml sterile phosphate-buffered saline, the solution will contain 1% human serum albumin (HSA) and 10% trehalose.
Reconstitution	It is recommended that 0.5 ml of sterile phosphate-buffered saline be added to the vial.
Storage	Lyophilized products should be stored at 2 to 8°C. Following reconstitution short-term storage at 4°C is recommended, and longer-term storage of aliquots at -18 to -20°C. Repeated freeze thawing is not recommended.
Activity	The ED ₅₀ of G-CSF is typically 0.01 - 0.03 ng/ml as measured in a cell proliferation assay using a murine myeloblastic M-NFS-60 cell line.

GENE INFORMATION

Gene Name	CSF3 colony stimulating factor 3 (granulocyte) [Homo sapiens]
Synonyms	CSF3; colony stimulating factor 3 (granulocyte); GCSF; G-CSF; MGC45931; colony stimulating factor 3; filgrastim; lenograstim; pluripoietin; granulocyte colony stimulating factor; Granulocyte colony-stimulating factor; OTTHUMP00000164327
Gene ID	1440

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_000759
Protein Refseq	NP_000750
UniProt ID	P09919
Chromosome Location	q11.2-q12
MIM	138970
Pathway	Cytokine-cytokine receptor interaction; opoietic cell lineage; Jak-STAT signaling pathway
Function	cytokine activity; enzyme binding; granulocyte colony-stimulating factor receptor binding; growth factor activity

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA