

Recombinant Human PDCD1LG2 protein, Avi & His-tagged, Biotinylated

Cat. No. PDCD1LG2-1281H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Human PD-L2 (recommended for biopanning) is expressed from human 293 cells (HEK293). It contains AA Leu 20 - Pro 219 (Accession # AAI13679). Biotinylation of this product is performed using Avitag technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin. The biotin to protein ratio is 0.5-1 as determined by the HABA assay.
Species	Human
Source	HEK293
ProteinLength	Leu 20 - Pro 219
Description	Programmed cell death 1 ligand 2 (PD-L2 or PDCD1 ligand 2) is also known as Butyrophilin B7-DC, CD antigen CD273, which belongs to the immunoglobulin superfamily or BTN/MOG family. The expression of PD-L2 is up-regulated by IFNG/IFN-gamma stimulation in monocytes and induced on dendritic cells grown from peripheral blood mononuclear cells with CSF2 and IL-4. PD-L2 Involved in the costimulatory signal, essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. PD-L2 interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.
Predicted N Terminal	Leu 20
Form	Lyophilized from 0.22

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	µm filtered solution in PBS, pH7.4, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 25.2 kDa. The protein migrates as 35-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 1.0 EU per µg by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20°C or lower. Please avoid repeated freeze-thaw cycles. No activity loss is observed after storage at: 4-8°C for 12 months in lyophilized state; -70°C for 3 months under sterile conditions after reconstitution
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	Biotin

GENE INFORMATION

Gene Name	PDCD1LG2 programmed cell death 1 ligand 2 [Homo sapiens (human)]
Official Symbol	PDCD1LG2
Synonyms	B7DC; Btdc; PDL2; CD273; PD-L2; PDCD1L2; bA574F11.2; B7 dendritic cell molecule; B7-DC; PD-1-ligand 2; PDCD1 ligand 2; butyrophilin B7-DC; programmed death ligand 2; programmed cell death 1 ligand 2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Gene ID 80380

mRNA Refseq NM_025239.3

Protein Refseq NP_079515.2

UniProt ID Q9BQ51

SDS-PAGE



Biotinylated Human PD-L2 (recommended for biopanning) on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA