

Recombinant Human MPL Protein, GST-tagged

Cat. No. MPL-5505H Lot. No. (See product label)

SPECIFICATION

Product Overview

Human MPL full-length ORF (NP_005364.1, 1 a.a. - 635 a.a.) recombinant protein with GST-tag at N-terminal.

Species

Human

Source

Wheat Germ

Description

In 1990 an oncogene, v-mpl, was identified from the murine myeloproliferative leukemia virus that was capable of immortalizing bone marrow hematopoietic cells from different lineages. In 1992 the human homologue, named, c-mpl, was cloned. Sequence data revealed that c-mpl encoded a protein that was homologous with members of the hematopoietic receptor superfamily. Presence of anti-sense oligodeoxynucleotides of c-mpl inhibited megakaryocyte colony formation. The ligand for c-mpl, thrombopoietin, was cloned in 1994. Thrombopoietin was shown to be the major regulator of megakaryocytopoiesis and platelet formation. The protein encoded by the c-mpl gene, CD110, is a 635 amino acid transmembrane domain, with two extracellular cytokine receptor domains and two intracellular cytokine receptor box motifs. TPO-R deficient mice were severely thrombocytopenic, emphasizing the important role of CD110 and thrombopoietin in megakaryocyte and platelet formation. Upon binding of thrombopoietin CD110 is dimerized and the JAK family of non-receptor tyrosine kinases, as well as the STAT family, the MAPK family, the adaptor protein Shc and the receptors themselves become tyrosine phosphorylated. [provided by RefSeq]

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	97.6 kDa
AA Sequence	MPSWALFMVTSCLLAPQNLAQVSSQDVSLASDSEPLKCFSRTFEDLTCFWDEEE AAPSGTYQLLYAYPREKPRACPLSSQSMPHFGTRYVCQFPDQEEVRLFFPLHLWVK NVFLNQTRTQRVLFVDSVGLPAPPSIIKAMGGSQPGELQISWEEPAPFISDFLRYELR YGPRDPKNSTGPTVIQLIATETCCPALQRPHSASALDQSPCAQPTMPWQDGPQKTS PSREASALTAEGGSCLISGLQPGNSYWLQLRSEPDGISLGGSWGWSLPTVDLPG DAVALGLQCFTLDLKNVTCQWQQQDHASSQGFFYHSRARCCPRDRYPIWENCEEE EKTNPGLQTPQFSRCHFCSRNDSEIHLVEVTTAPGTVHSYLGSPFWIHQAVRLPTPN LHWREISSGHLELEWQHPSSWAAQETCYQLRYTGEGHQDWKVLPPPLGARGGTL ELRPRSRYRLQLRARLNGPTYQGPWSSWSDPTRVETATETAWISLVTALHLVLGLS AVLGLLLLRLWQFPAHYRRLRHALWPSLPDLHRVLGQYLRDTAALSPPKATVSDTCE EVEPSLLEILPKSSERTPLPLCSSQAQMDYRRLQPSCLTGTMPLSVCPMAESGCC TTHIANHSYLPLSYWQQP
Applications	Enzyme-linked Immunoabsorbent Assay Western Blot (Recombinant protein) Antibody Production Protein Array
Notes	Best use within three months from the date of receipt of this protein.
Storage	Store at -80 centigrade. Aliquot to avoid repeated freezing and thawing.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
GENE INFORMATION	
Gene Name	MPL myeloproliferative leukemia virus oncogene [Homo sapiens]
Official Symbol	MPL

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	MPL; myeloproliferative leukemia virus oncogene; thrombopoietin receptor; CD110; TPOR; TPO-R; proto-oncogene c-Mpl; myeloproliferative leukemia protein; MPLV; C-MPL; THCYT2;
Gene ID	4352
mRNA Refseq	NM_005373
Protein Refseq	NP_005364
MIM	159530
UniProt ID	P40238
Quality Control Testing:	 12.5% SDS-PAGE Stained with Coomassie Blue.