

Active Recombinant Human Insulin-like Growth Factor Binding Protein 3

Cat. No. IGFBP3-145H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Insulin-like Growth Factor Binding Protein 3 DNA sequence encoding the human Insulin-like Growth Factor Binding Protein 3 sequence (containing the signal peptide sequence, and the mature human IGFBP-3 sequence) was expressed in modified <i>human 293 cells</i> .
Species	Human
Source	HEK293
Description	Insulin-like Growth Factor Binding Protein 3 is a member of the insulin-like growth factor binding protein (IGFBP) family. This family includes IGFBP-1 to IGFBP-6 that bind to the insulin-like growth factors (IGFs), thus regulating the half-life, activity, transport, and tissue distribution of the IGFs. Structurally, the IGFBP family of proteins is defined by highly homologous amino and carboxy terminal domains with a structurally diverse central region.
Amino Acid Sequence	GASSGGLGPVVRCEPCDARALAQCAPPAVCAELVREPGCGCCLTCALSEGQPCG IYTERCGSGLRCQPSPDEARPLQALLDGRGLCVNASAVSRLRAYLLPAPPAPGNAS ESEEDRSAGSVESPSVSSTHRVSDPKFHPLHSHKIIKKGHAKDSQRYKVDYESQSTD TQNFSSSESKRETEYGPCRREMEDTLNHLKFLNLSPRGVHIPNCDKKGFYKKKQCR PSKGRKRGFCWCVDKYGQPLPGYTTKGKEDVHCYSMQSK.
Molecular Mass	Insulin-like Growth Factor Binding Protein 3 migrates as a broad band between 40

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	and 45 kDa in SDS-PAGE due to post-translation modifications, in particular glycosylation. This compares with the unmodified IGFBP-3 that has a predicted molecular mass of 28.7 kDa.
pI	Insulin-like Growth Factor Binding Protein 3 separates into a number of isoforms with a pI between 6.8 and 8.2 in 2D PAGE due to post-translational modifications, in particular glycosylation. This compares with the unmodified IGFBP-3 that has a predicted pI of 8.76.
% Carbohydrate	Purified Insulin-like Growth Factor Binding Protein 3 consists of 25-35% carbohydrate by weight.
Purity	>95%, as determined by SDS-PAGE and visualized by silver stain.
Formulation	When reconstituted in 0.5 ml sterile phosphate-buffered saline, the solution will contain 1% human serum albumin (HSA) and 10% trehalose.
Reconstitution	It is recommended that 0.5 ml of sterile phosphate-buffered saline be added to the vial.
Storage	Lyophilized products should be stored at 2 to 8°C. Following reconstitution short-term storage at 4°C is recommended, and longer-term storage of aliquots at -18 to -20°C. Repeated freeze thawing is not recommended.
Activity	The ED50 of Insulin-like Growth Factor Binding Protein 3 is typically 0.13-0.20 µg/ml as measured by its ability to neutralize rhIGF-II mediated proliferation of the MCF-7 adenocarcinoma cell line.

GENE INFORMATION

Gene Name [IGFBP3 insulin-like growth factor binding protein 3 \[Homo sapiens \]](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	IGFBP3; insulin-like growth factor binding protein 3; IBP3; BP-53; binding protein 29; IGF-binding protein 3; growth hormone- dependent binding protein; acid stable subunit of the 140 K IGF complex
Gene ID	3486
mRNA Refseq	NM_000598
Protein Refseq	NP_000589
UniProt ID	P17936
Chromosome Location	7p13-p12
MIM	146732
Pathway	p53 signaling pathway; Diabetes pathways
Function	insulin-like growth factor I binding; metal ion binding; protein tyrosine phosphatase activator activity