

Recombinant Human Interleukin 1, Alpha, Fc Chimera

Cat. No. IL1A-109H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human interleukin 1, alpha encoding the signal peptide and extracellular domain of human IL-1 Receptor I (IL-1RI) (aa 1-333) , was expressed in modified <i>human 293 cells</i> . It was fused to the Fc region of human IgG1 (aa 93-330).
Species	Human
Source	HEK293
ProteinLength	1-333 a.a.
Description	Interleukin 1, alpha Fc Chimera is an 80kDa type I transmembrane glycoprotein expressed on T cells and fibroblasts. Structurally the extracellular domain (ECD) of IL-1R1 contains 3 Immunoglobulin like domains. Glycosylation of IL-1R1 has been shown to be necessary for optimal binding of IL-1, as blocking of glycosylation sites reduces binding to IL-1.
Amino Acid Sequence	LEADKCK EREEKIIL VS SANEIDVR PCP LNPNEHK GTITWYK DDSKTP VSTEQA SRIHQ HKEKLWF VPAKVED SGHYY CVVR NSSY CLRIK ISAK FVE NEP NL CY NAQAIFKQK LPVAGDGLVC PYMEFFK NENN EL PK LQ W YK DCKPLLLDNI HFSG VKDRLIVM NVAEKH RGNYTCH ASYTYLG KQYPITRVIE FIT LEE NKPT RPVIVSP ANETMEVDLG SGIQLICNVTGQL SDIAYWKWNGSVIDE DDPVLGE DYYSVE NPANKR RSTLI TVLNIS EIESRFYK HPFTCFANK THGIDAAYIQ LIYPVTNRSS NTKV DKK VEPKS CDKTH TCPPCP AP ELLGGPSV FL FPP KPK DTLMISRTPEVTCVVVDVSHEDPE VKFNWY VDGVEV HNAKTK PREEQY NSTY

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

RVVSVL TVLHQD WLNGK EYKCR VSNKALP APIEKTISKAKGQ PREPQ YTL
PPSRDE LTKNQVSLTCL VKGFYPSDIAV EWESNGQPEN NYKTTTPVLDSD GSF
FLYSKL TVDKS RWQQGNV FSCSVM HEALH NHYTQKSLSLSPGK.

Molecular Mass

Interleukin 1, alpha Fc Chimera migrates as a broad band between 65 and 100 kDa in SDS-PAGE due to post-translation modifications, in particular glycosylation. This compares with unmodified IL-1RI-Fc that has a predicted molecular mass of 63.6 kDa.

pI

Interleukin 1, alpha Fc Chimera separates into a number of isoforms with a pI between 5 and 8 in 2D PAGE due to post-translational modifications, in particular glycosylation. This compares with the unmodified IL-1RI-Fc that has a predicted pI of 7.

% Carbohydrate

Purified Interleukin 1, alpha Fc Chimera consists of 0-36% carbohydrate by weight.

Glycosylation

Interleukin 1, alpha Fc Chimera has N-linked and may have O-linked oligosaccharides.

Purity

>95%, as determined by SDS-PAGE and visualized by silver stain.

Formulation

When reconstituted in 0.5 ml sterile phosphate-buffered saline, the solution will contain 1% human serum albumin (HSA) and 10% trehalose.

Reconstitution

It is recommended that 0.5 ml of sterile phosphate-buffered saline be added to the vial.

Storage

Lyophilized products should be stored at 2 to 8°C. Following reconstitution short-term storage at 4°C is recommended, and longer-term storage of aliquots at -18 to -20°C. Repeated freeze thawing is not recommended.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GENE INFORMATION

Gene Name	IL1A interleukin 1, alpha [Homo sapiens]
Synonyms	IL1A; interleukin 1, alpha; IL1; IL-1A; IL1F1; IL1-ALPHA; preinterleukin 1 alpha; hematopoietin-1; pro-interleukin-1-alpha; Interleukin-1 alpha; IL-1 alpha; Hematopoietin-1; IL1A (IL1F1)
Gene ID	3552
mRNA Refseq	NM_000573
Protein Refseq	NP_000566
MIM	147760
UniProt ID	P01583
Chromosome Location	2q14
Pathway	Apoptosis; Cytokine-cytokine receptor interaction; Graft-versus-host disease; Hematopoietic cell lineage; MAPK signaling pathway; Prion diseases; Type I diabetes mellitus
Function	cytokine activity; interleukin-1 receptor binding; protein binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA