Xanthine Assay Kit

Cat. No. Kit-2476 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Xanthine is a purine base that can be found in most animal tissues and fluids. It is a product in the purine degradation pathway, produced by guanine deaminase from guanine, and by xanthine oxidoreductase from hypoxanthine. Xanthine is degraded to uric acid by xanthine oxidase. Clinically, xanthine and its derivatives act on sleep-inducing adenosine receptors as antagonists. Simple, direct and high-throughput assays for measuring xanthine find wide applications in research and drug discovery. xanthine assay kit uses a single Working Reagent that combines the xanthine oxidase reaction and color reaction in one step. The change in color intensity of the reaction product at 570 nm or fluorescence intensity at λex/em = 530/585 nm is directly proportional to xanthine concentration in the sample.
Storage	-20ºC
Shipping	On Ice
Size	100 tests
Detection method	OD570nm, FL530/585nm
Compatible Sample Types	Cell lysate, serum, and other biological samples.
Features & Benefits	Sensitive and accurate. Use as little as 10 µ samples. Linear detection range in 96-well plate for 30 minute incubation: 0.01 to 2. 5 mM xanthine for colorimetric assays and 3 to 250 uM for fluorimetric assays.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	Simple and convenient. The procedure involves addition of a single working reagent and incubation for 30 min at room temperature. Fast and high-throughput. Assays using 96-well plates and liquid handling system could allow simultaneous processing tens of thousands of samples per day.
Assay time	30 min
Sensitivity	FL 3 μM; OD 10 μM

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA