

Ketone Body Assay Kit

Cat. No. Kit-2506 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Ketone bodies (acetoacetic acid and 3-hydroxybutyric acid) are produced in the liver mainly from oxidation of fatty acids, and are normally present at low concentrations in urine and blood. Increased ketone concentrations in the blood may lead to metabolic acidosis, which has been associated with diabetes, childhood hypo-glycaemia, growth hormone deficiency, alcohol or salicylate intoxication and inborn errors of metabolism. Simple, direct and automation-ready procedures for measuring acetoacetic acid (AcAc) and 3-hydroxybutyric acid (BOH) are very desirable. ketone body assay is based on 3-hydroxybutyrate dehydrogenase catalyzed reactions, in which the change in NADH absorbance, measured at 340nm, is directly related to the AcAc and BOH concentrations.
Storage	-20°C
Shipping	On Ice
Size	200 tests
Detection method	OD340nm
Compatible Sample Types	Serum, plasma, urine etc
Features & Benefits	Sensitive and accurate. Uses 10 µL sample. Linear detection range of 0.12 to 8 mM for each ketone body in 96-well plate assay. Convenient. The procedure involves adding a single working reagent, and reading the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

optical density at room temperature.

High-throughput. Can be automated as a high-throughput 96-well plate assay for many samples per day.

Assay time

30 min

Sensitivity

0.12 mM

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA