

## Recombinant Human CD22 protein, Fc-tagged, APC labeled

**Cat. No.** CD22-562HA    **Lot. No.** (See product label)

### SPECIFICATION

|                         |                                                                                                                                                                                                                                                                                  |
|-------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product Overview</b> | Recombinant Human CD22 protein, Fc-tagged was Conjugated with NH2-Reactive APC(Allophycocyanin).                                                                                                                                                                                 |
| <b>Species</b>          | Human                                                                                                                                                                                                                                                                            |
| <b>Source</b>           | HEK293                                                                                                                                                                                                                                                                           |
| <b>Description</b>      | NH2-Reactive Allophycocyanin is used for antibody conjugation for immunostaining and cellular proteins for tracing. NH2-Reactive Allophycocyanin has succinimidyl ester groups, and can easily make a covalent bond with amino groups of protein without any activation process. |
| <b>Form</b>             | Liquid in sterile PBS buffer pH 7.4 (may contains tiny amount of reaction buffer and Allophycocyanin).                                                                                                                                                                           |
| <b>Endotoxin</b>        | <1EU/ug by LAL method.                                                                                                                                                                                                                                                           |
| <b>Stability</b>        | Sample is stable for at least 12 months when stored at -20 to -80°C from date of manufacture. For longer storage, add equal volume of glycerol to the sample solution and store at -20 to -80°C.                                                                                 |
| <b>Storage</b>          | Store under sterile conditions at -20 to -80°C immediately after receipt. It is recommended that the protein be aliquoted for optimal storage. Avoid freeze-thaw cycles.                                                                                                         |

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Related Product** CD22-562H**Conjugation** APC**GENE INFORMATION****Official Symbol** CD22**Synonyms**

CD22; CD22 molecule; CD22 antigen; B-cell receptor CD22; sialic acid binding Ig like lectin 2; SIGLEC 2; SIGLEC2; BL-CAM; T-cell surface antigen Leu-14; B-lymphocyte cell adhesion molecule; sialic acid binding Ig-like lectin 2; sialic acid-binding Ig-like lectin 2; SIGLEC-2; FLJ22814; MGC130020;

 Tel: 1-631-559-9269 1-516-512-3133 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA