

Recombinant Cynomolgus CXCL7 Protein, His-tagged

Cat. No. CXCL7-351C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Cynomolgus CXCL7 (XP_005555138.1) (Ser35-Asp128) was expressed with a polyhistidine tag at the C-terminus. Cynomolgus and Rhesus CXCL7 sequences are identical.
Species	Cynomolgus
Source	Yeast
ProteinLength	Ser35-Asp128
Description	Pro-platelet basic protein (PPBP) is also known as Chemokine (C-X-C motif) ligand 7 (CXCL7) and nucleosome assembly protein (Nap-2). Nap-2 / PPBP / CXCL7 is released in large amounts from platelets following their activation and is a platelet-derived growth factor that belongs to the CXC chemokine family. This growth factor is a potent chemoattractant and activator of neutrophils. Nap-2 / PPBP / CXCL7 has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells. Nap-2 is a ligand for CXCR1 and CXCR2, and Nap-2, Nap-2 (73), Nap-2 (74), Nap-2 (1-66), and most potent Nap-2 (1-63) are chemoattractants and activators for neutrophils.
Predicted N Terminal	Ser 35
Form	Lyophilized from sterile 30 % CAN, 0.1 % TFA. Normally 5 %- 8 % trehalose, mannitol

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

and 0.01% Tween80 are added as protectants before lyophilization.

Molecular Mass

The recombinant cynomolgus CXCL7 consists of 104 amino acids and predicts a molecular mass of 11.7 kDa.

Endotoxin

< 1.0 EU per µg protein as determined by the LAL method.

Purity

> 95 % as determined by SDS-PAGE.

Storage

Samples are stable for up to twelve months from date of receipt at -70 centigrade. Store it under sterile conditions at -20 centigrade to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Official Symbol

CXCL7

Synonyms

PPBP;platelet basic protein;Platelet basic protein;pro-platelet basic protein
(chemokine (C-X-C motif) ligand 7)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA