

Recombinant Cynomolgus IL20RB Protein, Fc-tagged

Cat. No. IL20RB-447C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Cynomolgus IL20RB (XP_005545913.1) (Asp30-Ala230) was expressed with the Fc region of human IgG1 at the C-terminus.
Species	Cynomolgus
Source	HEK293
ProteinLength	Asp30-Ala230
Description	IL20RB belongs to the type II cytokine receptor family. There are two kinds of type II cytokine receptors : cytokine receptors that bind type I and type II interferons; cytokine receptors that bind members of the interleukin-10 family (interleukin-10, interleukin-20 and interleukin-22). Type II cytokine receptors are similar to type I cytokine receptors except they do not possess the signature sequence WSXWS that is characteristic of type I receptors. They are expressed on the surface of certain cells, which bind and respond to a select group of cytokines. These receptors are related predominantly by sequence similarities in their extracellular portions that are composed of tandem Ig-like domains. The intracellular domain of type II cytokine receptors is typically associated with a tyrosine kinase belonging to the Janus kinase (JAK) family. IL20RB and IL20RA (MIM 605620) form a heterodimeric receptor for interleukin-20.
Predicted N Terminal	Asp 30
Form	Lyophilized from sterile PBS, pH 7.4. Normally 5 %- 8 % trehalose, mannitol and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	0.01% Tween80 are added as protectants before lyophilization.
Molecular Mass	The recombinant cynomolgus IL20RB consists of 439 amino acids and predicts a molecular mass of 49.2 kDa.
Endotoxin	< 1.0 EU per µg protein as determined by the LAL method.
Purity	> 90 % as determined by SDS-PAGE.
Storage	Samples are stable for up to twelve months from date of receipt at -70 centigrade. Store it under sterile conditions at -20 centigrade to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Official Symbol	IL20RB
Synonyms	IL20RB;interleukin 20 receptor subunit beta;interleukin-20 receptor subunit beta;interleukin 20 receptor beta

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA