

Recombinant Human CD80 protein, Mouse IgG2a Fc-tagged, low endotoxin

Cat. No. CD80-363H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Recombinant Human CD80 protein (Val35-Asn242)(low endotoxin), fused to Mouse IgG2a Fc tag at C-terminus, was expressed in human 293 cells (HEK293).

Species

Human

Source

HEK293

ProteinLength

208

Description

B7-1 and B7-2, together with their receptors CD28 and CTLA4, constitute one of the dominant co-stimulatory pathways that regulate T and Bcell responses. Although both CTLA4 and CD28 can bind to the same ligands, CTLA4 binds to B71 and B72 with a 20 100 fold higher affinity than CD28 and is involved in the downregulation of the immune response.

B-lymphocyte activation antigen B7-1 (referred to as B7) also known as cluster of Differentiation 80 (CD80), is a member of cell surface immunoglobulin superfamily and is expressed on activated B cells, activated T cells, macrophages and dendritic cells. It is the ligand for two different proteins on the T cell surface: CD28 (for autoregulation and intercellular association) and CTLA-4 (for attenuation of regulation and cellular disassociation). CD80 works in tandem with CD86 to prime T cells. CD80 plays a role in induction of innate immune responses by activating NF-κB-signaling pathway in macrophages. CD80 is thus regarded as promising therapeutic targets for autoimmune diseases and various carcinomas.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Lyophilized from 0.22 um filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 50.9 kDa. The protein migrates as 65-100 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 0.1 EU per ug by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 centigrade to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	CD80
Official Symbol	CD80
Synonyms	CD80; CD80 molecule; CD28LG, CD28LG1, CD80 antigen (CD28 antigen ligand 1, B7 1 antigen) , CD80 molecule; T-lymphocyte activation antigen CD80; B lymphocyte activation antigen B7; B7 1; B7.1; activation B7-1 antigen; costimulatory factor CD80; CTLA-4 counter-receptor B7.1; B-lymphocyte activation antigen B7; costimulatory molecule variant IgV-CD80; CD80 antigen (CD28 antigen ligand 1, B7-1 antigen); B7;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

BB1; B7-1; LAB7; CD28LG; CD28LG1

Gene ID 941


mRNA Refseq NM_005191

Protein Refseq NP_005182

MIM 112203

UniProt ID P33681

**Bioactivity-ELISA of
CD80-363H**




Immobilized Human CTLA-4, His Tag at 1 µg/mL (100 µL/well) can bind Human B7-1, Mouse IgG2a Fc Tag, low endotoxin with a linear range of 2-31 ng/mL (QC tested).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Bioactivity-ELISA of CD80-363H



Immobilized Human B7-1, Mouse IgG2a Fc Tag, low endotoxin at 5 µg/mL (100 µL/well) can bind Human PD-L1, Fc Tag with a linear range of 0.01-0.313 µg/mL (Routinely tested).

Bioactivity-ELISA of CD80-363H




Serial dilutions of Ipilimumab were added into Human B7-1, Mouse IgG2a Fc Tag, low endotoxin : Biotinylated Human CTLA-4, Fc,Avitag binding reactions. The half maximal inhibitory concentration (IC₅₀) is 0.8402 µg/mL (Routinely tested).

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127


45-1 Ramsey Road, Shirley, NY 11967, USA

Bioactivity-SPR of CD80-363H




Immobilized Human/Cynomolgus/Rhesus macaque CD28, Fc Tag (HPLC-verified) on CM5 Chip can bind Human B7-1, Mouse IgG2a Fc Tag, low endotoxin with an affinity constant of 2.72 μ M as determined in a SPR assay (Routinely tested).

Bioactivity-FACS of CD80-363H




Flow Cytometry assay shows that Human B7-1, Mouse IgG2a Fc Tag, low endotoxin can bind to 293 cell overexpressing human CTLA-4. The concentration of Human B7-1 is 1 μ g/mL (Routinely tested).

Bioactivity-FACS of CD80-363H



FACS analysis shows that the binding of Human B7-1, Mouse IgG2a Fc Tag, low endotoxin to 293 overexpressing CTLA-4 was inhibited by increasing concentration of neutralizing Anti-human CTLA-4 antibody. The concentration of Human B7-1 is 1 µg/mL. The IC₅₀ is 5.5 µg/mL (Routinely tested).

SDS-PAGE of CD80- 363H



Human B7-1, Mouse IgG2a Fc Tag, low endotoxin on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.