

Recombinant Mouse CD70 protein, Mouse IgG2a Fc-tagged, low endotoxin

Cat. No. CD70-330M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse CD70 protein (Gln47-Pro195)(low endotoxin), fused to Mouse IgG2a Fc tag at N-terminus, was expressed in human 293 cells (HEK293).
Species	Mouse
Source	HEK293
ProteinLength	149
Description	Cluster of Differentiation 70 (CD70) is also known as CD27 ligand (CD27L/CD27LG), TNFSF7, TNFSF7G, is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and is a surface antigen found on activated T-and B-lymphocytes and mature dendritic cells. Binding of CD70 to its receptor CD27 induces in priming, effector functions, differentiation and memory formation of T-cells, and thus is involved in the biological processes including T-cell activation, the proliferation of costimulated T-cells, as well as the generation of cytolytic T-cells. CD70 on T cells provides costimulatory signals that are required for T cell proliferation, clonal expansion and the promotion of effector T cell formation. CD70 on mouse B cell has been shown to inhibit terminal differentiation of activated B cells into plasma cells and enhances commitment to memory B cell responses. CD70 induces proliferation and IFN γ production, on NK cells.
Form	Lyophilized from 0.22 μ m filtered solution in 50 mM Tris, 100 mM Glycine, pH7.5,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	10% trehalose.
Molecular Mass	The protein has a calculated MW of 43.5 kDa. The protein migrates as 47-55 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 0.1 EU per ug by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 centigrade to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	CD70
Official Symbol	CD70
Synonyms	CD70; CD70 antigen; CD27-L; CD27 ligand; Ki-24 antigen; tumor necrosis factor ligand superfamily member 7; tumor necrosis factor (ligand) superfamily, member 7; Cd27l; CD27LG; Tnfsf7
Gene ID	21948

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq [NM_011617](#)

Protein Refseq [NP_035747](#)


UniProt ID [O55237](#)

Bioactivity of CD70-330M



Immobilized Human CD27 at 1 µg/mL (100 µL/well) can bind Mouse CD27 Ligand, Mouse IgG2a Fc Tag, low endotoxin with a linear range of 0.3-2 ng/mL (QC tested).

SDS-PAGE of CD70-330M



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Mouse CD27 Ligand, Mouse IgG2a Fc Tag, low endotoxin on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA