

Recombinant Human SIRPA protein, mouse IgG1 Fc-tagged (HPLC-verified)

Cat. No. SIRPA-346H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human SIRPA protein (Glu31-Arg370)(HPLC-verified), fused to Mouse IgG1 Fc tag at C-terminus, was expressed in human 293 cells (HEK293).
Species	Human
Source	HEK293
ProteinLength	340
Description	Tyrosine-protein phosphatase non-receptor type substrate 1 (SHPS1) is also known as CD172 antigen-like family member A (CD172a), Macrophage fusion receptor, MyD-1 antigen, Signal-regulatory protein alpha (SIRPA or SIRP alpha) or p84, is a member of the SIRP family, and also belongs to the immunoglobulin superfamily. SIRP alpha is Ubiquitous and highly expressed in brain. SIRPA/CD172a is immunoglobulin-like cell surface receptor for CD47 and acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRPA/SHPS-1 supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment and may play a key role in intracellular signaling during synaptogenesis and in synaptic function By similarity. SIRPA/MyD1 involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin and mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	production by mature dendritic cells.
Form	Lyophilized from 0.22 um filtered solution in 50 mM Tris, 100 mM Glycine, pH7.5, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 63.5 kDa. The predicted N-terminus is Glu 31. The reducing (R) protein migrates as 70-105 kDa in SDS-PAGE due to glycosylation.
Endotoxin	Less than 1.0 EU per ug by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 centigrade to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	SIRPA
Official Symbol	SIRPA
Synonyms	SIRPA; signal-regulatory protein alpha; protein tyrosine phosphatase, non receptor type substrate 1 , PTPNS1; tyrosine-protein phosphatase non-receptor type substrate 1; BIT; CD172a; MFR; MYD 1; P84; SHPS 1; SHPS1; SIRP; SIRP ALPHA 1;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SIRPalpha; SIRPalpha2; myd-1 antigen; inhibitory receptor SHPS-1; macrophage fusion receptor; CD172 antigen-like family member A; tyrosine phosphatase SHP substrate 1; brain-immunoglobulin-like molecule with tyrosine-based activation motifs; MYD-1; CD172A; PTPNS1

Gene ID 140885


mRNA Refseq NM_001040022

Protein Refseq NP_001035111

MIM 602461

UniProt ID P78324

Bioactivity-ELISA of SIRPA-346H




Immobilized Human CD47, His Tag at 2 µg/mL (100 µL/well) can bind Human SIRP alpha, Mouse IgG1 Fc Tag (HPLC-verified) with a linear range of 0.008-0.25 µg/mL (QC tested).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Bioactivity-ELISA of SIRPA-346H




Serial dilutions of Anti-Human CD47 Neutralizing Antibody were added into Human SIRP alpha, Mouse IgG1 Fc Tag (HPLC-verified): Biotinylated Human CD47, Fc,Avitag binding reactions. The half maximal inhibitory concentration (IC₅₀) is 0.2006 µg/mL (Routinely tested).

Bioactivity-FACS of SIRPA-346H




FACS assay shows that Human SIRP alpha, Mouse IgG1 Fc Tag (HPLC-verified) can bind to Jurkat cell expressing CD47. The concentration of SIRP alpha used is 0.3 µg/mL (Routinely tested).

Bioactivity-FACS of SIRPA-346H



FACS analysis shows that the binding of Human SIRP alpha, Mouse IgG1 Fc Tag (HPLC-verified) to Jurkat expressing CD47 was inhibited by increasing concentration of neutralizing anti-CD47 antibody. The concentration of SIRP alpha used is 0.3 µg/mL. IC₅₀=0.1318 µg/mL (Routinely tested).


SDS-PAGE of SIRPA- 346H



Human SIRP alpha, Mouse IgG1 Fc Tag (HPLC-verified) on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The

purity of the protein is greater than 95%.

SEC-HPLC of SIRPA-346H



The purity of Human SIRP alpha, Mouse IgG1 Fc Tag (HPLC-verified) was greater than 90% as determined by SEC-HPLC.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA