

Recombinant Human TNFRSF13B protein, Fc-tagged

Cat. No. TNFRSF13B-359H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human TNFRSF13B protein (Ser2-Thr166), fused to human IgG1 Fc tag at C-terminus, was expressed in human 293 cells (HEK293).
Species	Human
Source	HEK293
ProteinLength	165
Description	Transmembrane activator and CAML interactor (TACI), also known as tumor necrosis factor receptor superfamily member 13B (TNFRSF13B). It was originally discovered because of its ability to interact with calcium-modulator and cyclophilin ligand (CAML). TACI was later found to play a crucial role in humoral immunity by interacting with two members of the TNF family: BAFF and APRIL. The present study demonstrated that, in NSCLC, a proliferation-inducing ligand (APRIL), B-cell maturation antigen (BCMA) and transmembrane activator and CAML interactor (TACI) proteins are abnormally expressed by immunohistochemistry, reverse transcription-quantitative polymerase chain reaction and western blotting. In addition, the expression of APRIL, BCMA and TACI were observed to be involved in extracellular signal-regulated kinase (ERK)1/2 activation in A549 cells.
Form	Lyophilized from 0.22 um filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 44.6 kDa. The protein migrates as 36 kDa, 37

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	kDa and 46 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 1.0 EU per ug by the LAL method.
Purity	>90% as determined by reduced SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 centigrade to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	TNFRSF13B
Official Symbol	TNFRSF13B
Synonyms	TNFRSF13B; tumor necrosis factor receptor superfamily, member 13B; tumor necrosis factor receptor superfamily member 13B; CD267; TACI; tumor necrosis factor receptor 13B; transmembrane activator and CAML interactor; CVID; CVID2; TNFRSF14B; FLJ39942; MGC39952; MGC133214
Gene ID	23495
mRNA Refseq	NM_012452

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_036584

MIM 604907


UniProt ID O14836

**Bioactivity-ELISA of
TNFRSF13B-359H**



Immobilized Human BAFF, His Tag at 2 µg/mL (100 µL/well) can bind Human TACI/TNFRSF13B, Fc Tag with a linear range of 2-16 ng/mL (QC tested).

**SDS-PAGE of
TNFRSF13B-359H**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Human TACI/TNFRSF13B, Fc Tag on SDS-PAGE under reducing (R) condition.
The gel was stained overnight with Coomassie Blue. The purity of the protein is
greater than 90%.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA