

## Active Recombinant Bovine CXCL5 protein

**Cat. No.** CXCL5-42B    **Lot. No.** (See product label)

### SPECIFICATION

|                         |                                                                                                                                                                                      |
|-------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product Overview</b> | Active Recombinant Bovine CXCL5 protein(P80221)(37-112 aa) was expressed in E.coli.                                                                                                  |
| <b>Species</b>          | Bovine                                                                                                                                                                               |
| <b>Source</b>           | E.coli                                                                                                                                                                               |
| <b>ProteinLength</b>    | 37-112 aa                                                                                                                                                                            |
| <b>Form</b>             | Lyophilized from a 0.2 µ filtered concentrated solution in 20 mM PB, 500 mM NaCl, pH 7.0.                                                                                            |
| <b>Bio-activity</b>     | Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human neutrophils is in a concentration range of 10-50 ng/ml. |
| <b>Molecular Mass</b>   | Approximately 8.0 kDa, a single non-glycosylated polypeptide chain containing 76 amino acids.                                                                                        |
| <b>AASequence</b>       | GPVAAVVREL RCVCLTTTPG IHPKTVSDLQ VIAAGPQCSK VEVIATLKNG<br>REVCLDPEAP LIKKIVQKIL DSGKNN                                                                                               |
| <b>Endotoxin</b>        | Less than 0.1 EU/g of rBoGCP-2/CXCL6 as determined by LAL method.                                                                                                                    |
| <b>Purity</b>           | >97% by SDS-PAGE and HPLC analyses.                                                                                                                                                  |

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

|                       |                                                                                                                                                                                                                                                                                                                                                                                           |
|-----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Storage</b>        | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. - 12 months from date of receipt, -20 to -70 °C as supplied. - 1 month, 2 to 8 °C under sterile conditions after reconstitution. - 3 months, -20 to -70 °C under sterile conditions after reconstitution.                                                                                                             |
| <b>Reconstitution</b> | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq -20$ °C. Further dilutions should be made in appropriate buffered solutions. |

 Tel: 1-631-559-9269 1-516-512-3133 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA